arXiv:1505.07536v1 [math.SP] 28 May 2015

Continuous Dependence of the n-th Eigenvalue on Self-adjoint Discrete Sturm-Liouville Problem

Hao Zhu, Yuming Shi [†]

Department of Mathematics, Shandong University
Jinan, Shandong 250100, P. R. China

Abstract. This paper is concerned with continuous dependence of the n-th eigenvalue on self-adjoint discrete Sturm-Liouville problems. The n-th eigenvalue is considered as a function in the space of the problems. A necessary and sufficient condition for all the eigenvalue functions to be continuous and several properties of the eigenvalue functions in a set of the space of the problems are given. They play an important role in the study of continuous dependence of the n-th eigenvalue function on the problems. Continuous dependence of the n-th eigenvalue function on the equations and on the boundary conditions is studied separately. Consequently, the continuity and discontinuity of the n-th eigenvalue function are completely characterized in the whole space of the problems. Especially, asymptotic behaviors of the n-th eigenvalue function near each discontinuity point are given.

Keywords: self-adjoint discrete Sturm-Liouville problem; eigenvalue function; continuous dependence; asymptotic behavior.

2000 *Mathematics Subject Classification*: 39A12; 34B24; 39A70.

1. Introduction

A self-adjoint discrete Sturm-Liouville problem (briefly, SLP) considered in the present paper consists of a symmetric discrete Sturm-Liouville equation (briefly, SLE)

$$-\nabla(f_n\Delta y_n)+q_ny_n=\lambda w_ny_n,\qquad n\in[1,N],\tag{1.1}$$

and a self-adjoint boundary condition (briefly, BC)

$$A\begin{pmatrix} y_0\\ f_0\triangle y_0\end{pmatrix}+B\begin{pmatrix} y_N\\ f_N\triangle y_N\end{pmatrix}=0,\tag{1.2}$$

where $N\geq 2$ is an integer, Δ and ∇ are the forward and backward difference operators, respectively, i.e., $\Delta y_n=y_{n+1}-y_n$ and $\nabla y_n=y_n-y_{n-1}$; $f=\{f_n\}_{n=0}^N$, $q=\{q_n\}_{n=1}^N$ and $w=\{w_n\}_{n=1}^N$ are real-valued sequences such that

---

[†] The corresponding author.

Email addresses: haozhusdu@163.com(H. Zhu), ymshi@sdu.edu.cn(Y. Shi).

$$f_n \neq 0 \ \text{ for } \ n \in [0, N], \quad w_n > 0 \ \text{ for } \ n \in [1, N]; \tag{1.3}$$

λ is the spectral parameter; the interval $[M, N]$ denotes the set of integers $\{M, M+1, \cdots, N\}$; A and B are 2×2 complex matrices such that rank(A,B)=2, and satisfy the following self-adjoint boundary condition:

$$A \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} A^* = B \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} B^*, \tag{1.4}$$

while A^* denotes the complex conjugate transpose of A.

Throughout this paper, by $\mathbb{C}$, $\mathbb{R}$, and $\mathbb{N}$ denote the sets of the complex, real, and natural numbers, respectively; and by $\bar{z}$ denotes the complex conjugate of $z \in \mathbb{C}$. When a capital Latin letter stands for a matrix, its entries are denoted by the corresponding lower case letter with two indices. For example, the entries of a matrix C are c_{ij}'s.

The SLP (1.1)–(1.2) has attracted increasing attention by many scholars, for its applications in various areas of physical science [1, 9]. The n-th eigenvalue of the SLP always stands for an important value in the physical problems and it varies as the SLP varies. A natural question is how it varies as the SLP varies. In the present paper, we are interested in continuous dependence of the n-th eigenvalue on the SLP. We shall consider the n-th eigenvalue as a function in the space of the SLPs, and mainly study its continuity and discontinuity, and characterize its asymptotic behaviors near each discontinuity point.

The continuous dependence of the n-th eigenvalue on self-adjoint continuous Sturm-Liouville problems has been studied quite deeply and some elegant results have been obtained (cf., [6, 8, 10, 11, 14, 21]). Now, we shall briefly recall some existing results of continuous dependence of the n-th eigenvalue on self-adjoint continuous SLPs. A self-adjoint continuous SLP consists of a differential SLE

$$-(p(t)y')' + q(t)y = \lambda w(t)y, \qquad t \in (a, b), \tag{1.5}$$

and a BC

$$A \begin{pmatrix} y(a) \\ (py')(a) \end{pmatrix} + B \begin{pmatrix} y(b) \\ (py')(b) \end{pmatrix} = 0, \tag{1.6}$$

where $-\infty < a < b < +\infty$; $1/p, q, w \in L((a, b), \mathbb{R})$ with $p, w > 0$ almost everywhere in (a, b), while $L((a, b), \mathbb{R})$ denotes the space of Lebesgue integrable real functions in (a, b); A and B are 2×2 complex matrices such that rank(A,B)=2 and (1.4) holds. It is well-known that the problem (1.5)–(1.6) has infinitely countable eigenvalues, which are all real and can be arranged in the following non-decreasing order:

$$\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq \cdots \leq \lambda_n \leq \cdots \tag{1.7}$$

with $\lambda_n \to +\infty$ as $n \to +\infty$, counting repeatedly according to their multiplicities. Using the variational method, Courant and Hilbert showed that the n-th eigenvalue λ_n is continuously dependent on the problem under the assumptions that the coefficient functions p,

q and w in (1.5) are continuous functions in (a, b) and the BC (1.6) is of a special class [6]. In 1997, Everitt, Möller, and Zettl showed that the n-th eigenvalue λ_n does not depend continuously on the BCs in general [8]. Later, Kong, Wu, and Zettl deeply studied this problem in 1999 [10]. They showed that the n-th eigenvalue λ_n depends continuously on the SLEs, found its discontinuity set of the space of BCs

$$\begin{aligned}\mathcal{J}^{\mathbb{C}} = & \left\{[e^{i\theta}K \mid -I] : K \in SL(2,\mathbb{R}), k_{12} = 0, \theta \in [0,\pi)\right\} \\ & \bigcup \left\{ \begin{bmatrix} a_1 & a_2 & 0 & 0 \\ 0 & 0 & b_1 & b_2 \end{bmatrix} : a_2 b_2 = 0 \right\},\end{aligned} \tag{1.8}$$

and gave its asymptotic behaviors near each discontinuity point.

The spectral theory of self-adjoint discrete Sturm-Liouville problems has attracted a great deal of interest from many authors and some good results have been obtained (cf., [1, 3–5, 9, 12, 15–20, 22]). In [17], the second author of the present paper with her coauthor Chen studied some regular self-adjoint spectral problems for second-order vector difference equations, which include (1.1)–(1.2), and gave several spectral results, including the reality of the eigenvalues, the finiteness of the number of the eigenvalues, and a formula for counting the number of the eigenvalues. Based on these results, the problem (1.1)–(1.2) has k eigenvalues (counting multiplicities), which are real and can be arranged in the following non-decreasing order:

$$\lambda_0 \le \lambda_1 \le \lambda_2 \le \cdots \le \lambda_{k-1},$$

where k can be determined (see Lemma 2.4). Note that the analytic and geometric multiplicities of an eigenvalue of an SLP (1.1)–(1.2) are the same [18, 22]. Recently, we studied some problems about dependence of the eigenvalues of (1.1)–(1.2) on the problems in [22]. We gave the topologies and geometric structures of the space of the SLPs (1.1)–(1.2), showed that each eigenvalue of a given SLP lives in one or two continuous eigenvalue branches, and studied analyticity, differentiability and monotonicity of continuous eigenvalue branches.

It is evident that the n-th eigenvalue depends on the SLP (1.1)–(1.2) and then can be regarded as a function in the space of the SLPs. So, based on the work in [22], we shall mainly study continuous dependence of the n-th eigenvalue on the problem in the present paper. We shall give out its continuity and discontinuity sets in the space of the SLPs (1.1)–(1.2), and characterize its asymptotic behaviors near each discontinuity point.

From Example 5.3 in [22], we have found that the index of eigenvalues in a continuous eigenvalue branch may change as the problem varies. This may lead the discontinuity of the n-th eigenvalue function. For convenience, we shall briefly recall this example.

Example 1.1 [22, Example 5.3]. Consider (1.1)–(1.2), where

$$N=2,\ f_0=f_1=f_2=1,\ q_1=q_2=0,\ w_1=w_2=1,$$

and

$$A=A(\alpha)=\begin{pmatrix}\cos\alpha & -\sin\alpha\\ 0 & 0\end{pmatrix},\ B=\begin{pmatrix}0 & 0\\ 0 & -1\end{pmatrix}.$$

We have showed that the SLP with $\alpha=3\pi/4$ has exactly one eigenvalue $\lambda_0=1$, and each SLP with $\alpha\in[0,3\pi/4)\cup(3\pi/4,\pi)$ has exactly the following two eigenvalues:

$$\lambda_0(\alpha)=\begin{cases}\lambda_-(\alpha) & \text{if } \alpha\in[0,3\pi/4),\\ \lambda_+(\alpha) & \text{if } \alpha\in(3\pi/4,\pi),\end{cases}\qquad \lambda_1(\alpha)=\begin{cases}\lambda_+(\alpha) & \text{if } \alpha\in[0,3\pi/4),\\ \lambda_-(\alpha) & \text{if } \alpha\in(3\pi/4,\pi),\end{cases}$$

where

$$\lambda_\pm(\alpha)=\frac{3\cos\alpha+2\sin\alpha\pm\sqrt{\cos^2\alpha+4\sin(2\alpha)+4}}{2(\cos\alpha+\sin\alpha)}.$$

So, there are exactly the following three continuous eigenvalue branches:

$$\begin{aligned}\Lambda_1(\alpha)\ &=\lambda_0(\alpha)\ \text{ for } \alpha\in[0,3\pi/4),\\ \Lambda_2(\alpha)\ &=\begin{cases}\lambda_1(\alpha) & \text{if } \alpha\in[0,3\pi/4),\\ 1 & \text{if } \alpha=3\pi/4,\\ \lambda_0(\alpha) & \text{if } \alpha\in(3\pi/4,\pi),\end{cases}\\ \Lambda_3(\alpha)\ &=\lambda_1(\alpha)\ \text{ for } \alpha\in(3\pi/4,\pi).\end{aligned}$$

See Figure 1.1.



Figure 1.1.

As functions in $\alpha\in[0,\pi)$, the eigenvalues λ_0 and λ_1 are not continuous at $\alpha=3\pi/4$, and have the following asymptotic behaviors near $3\pi/4$:

$$\lim_{\alpha\to3\pi/4^-}\lambda_0(\alpha)=-\infty,\quad \lim_{\alpha\to3\pi/4^-}\lambda_1(\alpha)=\lambda_0(3\pi/4),$$
$$\lim_{\alpha\to3\pi/4^+}\lambda_0(\alpha)=\lambda_0(3\pi/4),\quad \lim_{\alpha\to3\pi/4^+}\lambda_1(\alpha)=+\infty.$$

Let $\mathcal{O}$ be a set in the space of the SLPs (1.1)–(1.2). Through the above observation, in the study of the continuity of the n-th eigenvalue function restricted in $\mathcal{O}$, it seems very important whether the number of eigenvalues of each SLP in $\mathcal{O}$ is equal or not. In fact, we shall show that all the eigenvalue functions restricted in $\mathcal{O}$ are continuous if and only if the number of eigenvalues of each SLP in $\mathcal{O}$ is equal (see Theorem 2.1).

Our study in the present paper was inspired by the remarkable work in [10] for the continuous case. Note that the number of the eigenvalues is finite for the discrete problem (1.1)–(1.2) and infinite for the continuous problem (1.5)–(1.6). This difference will result in some differences between properties of their eigenvalue functions. Consequently, our method used in the present paper is quite different from that used in the continuous case. We shall list six aspects on these differences as follows. **Firstly**, it was shown in [10] that if the minimal eigenvalue function λ_0 is bounded from below in a set of the space of the SLPs (1.5)–(1.6), then the n-th eigenvalue function restricted in the set is continuous for each $n \geq 0$. However, a similar conclusion is not true in the discrete case (see Example 2.1). Instead, as we have remarked in the above, the continuity of the eigenvalue functions in a set can be completely determined by the number of eigenvalues of each SLP in the set (see Theorem 2.1). **Secondly**, unlike that in the continuous case, the n-th eigenvalue function is not continuously dependent on the SLE (1.1) in general in the discrete case. **Thirdly**, the discontinuity set of the n-th eigenvalue function in the space of BCs in the continuous case is different from that in the discrete case (see (1.8) and (4.12)). They may be identified in a certain sense by letting $f_0 \to +\infty$ in the discrete case. **Fourthly**, the authors in [10] employed the Prüfer transformation of (1.5) and some inequalities among eigenvalues of SLPs (1.5)–(1.6) given in [7] in their discussions. Though a discrete Prüfer transformation was established in [2] and several inequalities among eigenvalues of discrete SLPs were obtained in [19, 20], we have found that it is quite difficult for us to employ a similar method to study this discrete problem. Instead, we shall directly study several properties of the eigenvalue functions and make use of some spectral results of second-order difference equations given in [17, 22]. In order to study asymptotic behaviors of the n-th eigenvalue function λ_n near a discontinuity point in the space of the SLPs (1.1)–(1.2), we shall first study asymptotic behaviors of λ_n in a certain direction near the discontinuity point. This way is shown to be convenient for our study in the discrete case. **Fifthly**, continuous dependence of the eigenvalue functions on the BCs is investigated via the local coordinate systems in the space of BCs (1.2) directly, instead of being divided into the separated and coupled cases. **Finally**, asymptotic behaviors of the n-th eigenvalue function near a discontinuity point in the space of BCs (1.2) are more complicated than those in the continuous case.

The rest of this paper is organized as follows. In Section 2, some notations and lemmas are introduced. A necessary and sufficient condition for all the eigenvalue functions to be continuous and several properties of the eigenvalue functions in a set of the space of SLPs are given. In Section 3, continuous dependence of the n-th eigenvalue function on the SLE is completely characterized for a fixed BC. In Section 4, continuous dependence of the n-th eigenvalue function on the BC is completely characterized for a fixed SLE. In Section 5, continuous dependence of the n-th eigenvalue function on the SLP is studied. Then its continuity and discontinuity in the space of the SLPs are completely characterized.

Remark 1.1. We shall apply some results obtained in the present paper to study inequalities among the eigenvalues of general self-adjoint SLPs in our forthcoming paper.

2. Preliminaries

In this section, some notations and lemmas are introduced. This section is divided into three parts. In Section 2.1, the description of the space of the SLPs is introduced. Section 2.2 collects some basic properties of eigenvalues of the SLPs. In Section 2.3, a necessary and sufficient condition for all the eigenvalue functions to be continuous and several properties of the eigenvalue functions in a set of the space of the SLPs are given. They are useful in the sequent sections.

2.1. Space of self-adjoint discrete SLPs

In order to discuss continuous dependence of the n-th eigenvalue on the SLP, we need to know how to measure the closeness of two SLEs and of two BCs.

Let the SLE (1.1) be abbreviated as $(1/f, q, w)$. Then the space of the SLEs can be written as

$$\Omega_N^{\mathbb{R},+} := \{(1/f, q, w) \in \mathbb{R}^{3N+1} : (1.3) \text{ holds}\},$$

and is equipped with the topology deduced from the real space $\mathbb{R}^{3N+1}$. Note that $\Omega_N^{\mathbb{R},+}$ has 2^{N+1} connected components. Bold faced lower case Greek letters, such as $\boldsymbol{\omega}$, are used to denote elements of $\Omega_N^{\mathbb{R},+}$. For convenience, the maximum norm in $\mathbb{R}^{3N+1}$ will be used:

$$\|(1/f, q, w)\| = \max\left\{|1/f_0|, \max_{1\le n\le N}\{|1/f_n|, |q_n|, |w_n|\}\right\}.$$

The quotient space

$$\mathcal{A}^{\mathbb{C}} := \mathrm{M}^*_{2,4}(\mathbb{C})/\mathrm{GL}(2,\mathbb{C}),$$

equipped with the quotient topology, is taken as the space of general BCs; that is, each BC is an equivalence class of coefficient matrices of system (1.2), where

$$M^*_{2,4}(\mathbb{C}) := \{2 \times 4 \text{ complex matrix } (\mathrm{A}, \mathrm{B}) : \mathrm{rank}(A, B) = 2\},$$
$$GL(2,\mathbb{C}) := \{2 \times 2 \text{ comlplex matrix } T : \det T \neq 0\}.$$

The BC represented by (1.2) is denoted by $[A \,|\, B]$. Bold faced capital Latin letters, such as $\mathbf{A}$, are also used for BCs. The space of self-adjoint BCs is denoted by $\mathcal{B}^{\mathbb{C}}$. The following result gives the topology and geometric structure of $\mathcal{B}^{\mathbb{C}}$.

Lemma 2.1 [22, Theorem 2.2]. *The space $\mathcal{B}^{\mathbb{C}}$ equals the union of the following relative open sets:*

$$\begin{aligned}
\mathcal{O}^{\mathbb{C}}_{1,3} &= \left\{ \left[\begin{array}{cccc} 1 & a_{12} & 0 & \bar{z} \\ 0 & z & -1 & b_{22} \end{array} \right] : \ a_{12}, b_{22} \in \mathbb{R}, z \in \mathbb{C} \right\}, \\
\mathcal{O}^{\mathbb{C}}_{1,4} &= \left\{ \left[\begin{array}{cccc} 1 & a_{12} & \bar{z} & 0 \\ 0 & z & b_{21} & 1 \end{array} \right] : \ a_{12}, b_{21} \in \mathbb{R}, z \in \mathbb{C} \right\}, \\
\mathcal{O}^{\mathbb{C}}_{2,3} &= \left\{ \left[\begin{array}{cccc} a_{11} & -1 & 0 & \bar{z} \\ z & 0 & -1 & b_{22} \end{array} \right] : \ a_{11}, b_{22} \in \mathbb{R}, z \in \mathbb{C} \right\}, \\
\mathcal{O}^{\mathbb{C}}_{2,4} &= \left\{ \left[\begin{array}{cccc} a_{11} & -1 & \bar{z} & 0 \\ z & 0 & b_{21} & 1 \end{array} \right] : \ a_{11}, b_{21} \in \mathbb{R}, z \in \mathbb{C} \right\}.
\end{aligned} \tag{2.1}$$

Moreover, $\mathcal{B}^{\mathbb{C}}$ is a connected and compact real-analytic manifold of dimension 4.

Lemma 2.1 says that $\mathcal{O}^{\mathbb{C}}_{1,3}$, $\mathcal{O}^{\mathbb{C}}_{1,4}$, $\mathcal{O}^{\mathbb{C}}_{2,3}$, and $\mathcal{O}^{\mathbb{C}}_{2,4}$ together form an atlas of local coordinate systems in $\mathcal{B}^{\mathbb{C}}$. If $\mathbf{A} \in \mathcal{O}^{\mathbb{C}}_{1,3}$, then the corresponding coordinate chart is given by

$$\hat{\varphi} : \left[\begin{array}{cccc} 1 & a_{12} & 0 & \bar{z} \\ 0 & z & -1 & b_{22} \end{array} \right] \to (a_{12}, a, b, b_{22}),$$

where $z = a + ib$, a, $b \in \mathbb{R}$. Others are given similarly. Thus, each of the four open sets in (2.1) can be identified with $\mathbb{R}^4$.

The space $\Omega^{\mathbb{R},+}_N \times \mathcal{B}^{\mathbb{C}}$ of the SLPs is a real-analytic manifold of dimension $3N + 5$ and has 2^{N+1} connected components.

The following result gives the canonical forms of separated and coupled self-adjoint BCs, respectively.

Lemma 2.2 [21, Theorem 10.4.3]. *Each separated self-adjoint BC can be written as*

$$\mathbf{S}_{\alpha,\beta} := \left[\begin{array}{cccc} \cos\alpha & -\sin\alpha & 0 & 0 \\ 0 & 0 & \cos\beta & -\sin\beta \end{array} \right], \tag{2.2}$$

where

$$\alpha \in [0, \pi), \beta \in (0, \pi];$$

and each coupled self-adjoint BC can be written as

$$[e^{i\gamma} K \,|\, -I],$$

where

$$\gamma \in [0, \pi), \quad K \in SL(2, \mathbb{R}) := \{2 \times 2 \textit{ real matrix } M : \det M = 1\}.$$

The spaces of separated and coupled self-adjoint BCs are denoted by $\mathcal{B}_S$ and $\mathcal{B}_C$, respectively. So $\mathcal{B}^{\mathbb{C}} = \mathcal{B}_S \cup \mathcal{B}_C$.

2.2. Basic properties of eigenvalues

In this subsection, some basic properties of eigenvalues of the SLPs are introduced.

For each $\lambda \in \mathbb{C}$, let $\phi(\lambda)$ and $\psi(\lambda)$ be the solutions of (1.1) satisfying the following initial conditions:

$$\phi_0(\lambda) = 1, f_0\Delta\phi_0(\lambda) = 0; \ \ \psi_0(\lambda) = 0, \ f_0\Delta\psi_0(\lambda) = 1,$$

separately. It follows from [22] that the leading terms of $\phi_N(\lambda)$, $\psi_N(\lambda)$, $f_N\Delta\phi_N(\lambda)$, and $f_N\Delta\psi_N(\lambda)$ as polynomials of λ are

$$\begin{aligned} &(-1)^{N-1}\left(\prod_{i=1}^{N-1}(w_i/f_i)\right)\lambda^{N-1}, \quad (-1)^{N-1}\left((1/f_0)\prod_{i=1}^{N-1}(w_i/f_i)\right)\lambda^{N-1}, \\ &(-1)^{N}\left(w_N\prod_{i=1}^{N-1}(w_i/f_i)\right)\lambda^{N}, \quad (-1)^{N}\left((w_N/f_0)\prod_{i=1}^{N-1}(w_i/f_i)\right)\lambda^{N}, \end{aligned} \tag{2.3}$$

respectively.

Lemma 2.3 [22, Lemmas 3.2 and 3.3]. *A number $\lambda \in \mathbb{R}$ is an eigenvalue of the SLP* (1.1)-(1.2) *if and only if λ is a zero of the characteristic polynomial*

$$\Gamma(\lambda) = \det A + \det B + G(\lambda),$$

where

$$G(\lambda) := c_{11}\phi_N(\lambda) + c_{12}\psi_N(\lambda) + c_{21}f_N\Delta\phi_N(\lambda) + c_{22}f_N\Delta\psi_N(\lambda),$$

$$C := \begin{pmatrix} b_{11} & b_{21} \\ b_{12} & b_{22} \end{pmatrix}\begin{pmatrix} a_{22} & -a_{21} \\ -a_{12} & a_{11} \end{pmatrix}.$$

When we count the eigenvalues of an SLP in a domain in $\mathbb{R}$, their multiplicities will be taken into account.

Let $(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$. Set

$$r = r(\boldsymbol{\omega}, \mathbf{A}) := \operatorname{rank}\begin{pmatrix} -a_{11} + f_0a_{12} & b_{12} \\ -a_{21} + f_0a_{22} & b_{22} \end{pmatrix}. \tag{2.4}$$

Obviously, $0 \le r \le 2$.

Lemma 2.4 [22, Lemma 3.4]. *The number of eigenvalues of $(\boldsymbol{\omega}, \mathbf{A})$ is equal to $N - 2 + r$, where r is defined by* (2.4).

The above results can be deduced from [17, Theorem 4.1].

By (2.3) and Lemma 2.3, the coefficient of λ^N in the polynomial $\Gamma(\lambda)$ is

$$\theta(\boldsymbol{\omega}, \mathbf{A}) := (-1)^N\left(w_N\prod_{i=1}^{N-1}(w_i/f_i)\right)\left[(a_{11}b_{22} - a_{21}b_{12})/f_0 + a_{22}b_{12} - a_{12}b_{22}\right]. \tag{2.5}$$

Thus, by Lemma 2.4, there are exactly the following three cases:

(i) $(\boldsymbol{\omega}, \mathbf{A})$ has exactly N eigenvalues in the case that $\theta(\boldsymbol{\omega}, \mathbf{A}) \neq 0$;

(ii) $(\boldsymbol{\omega}, \mathbf{A})$ has exactly $N-1$ eigenvalues in the case that $\theta(\boldsymbol{\omega}, \mathbf{A}) = 0$ and $\mathbf{A} \neq \mathbf{A}(1/f_0)$, where

$$\mathbf{A}(1/f_0) = \begin{bmatrix} 1 & 1/f_0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix};$$

(iii) $(\boldsymbol{\omega}, \mathbf{A})$ has exactly $N-2$ eigenvalues in the case that $\theta(\boldsymbol{\omega}, \mathbf{A}) = 0$ and $\mathbf{A} = \mathbf{A}(1/f_0)$.

By $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ denotes the n-th eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$. When the SLE is fixed, $\lambda_n(\mathbf{A})$ is also used for $\mathbf{A} \in \mathcal{B}^{\mathbb{C}}$; when the BC is fixed, $\lambda_n(\boldsymbol{\omega})$ is also used for $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+}$, and etc.

The following result is a generation of [22, Corollary 3.3].

Lemma 2.5. *Assume that $\mathcal{O}$ is a set of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$, and $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{O}$. Let r_1 and r_2 be two real numbers with $r_1 < r_2$ such that neither of them is an eigenvalue of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$, and $n \geq 0$ be the number of eigenvalues of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in the interval $[r_1, r_2]$. Then there exists a neighborhood $\mathcal{U}$ of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in $\mathcal{O}$ such that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{U}$ has exactly n eigenvalues in $[r_1, r_2]$, which all lie in (r_1, r_2).*

Proof. Since the proof is similar to that of [22, Theorem 3.4], we omit its details.

Lemma 2.6 [22, Theorem 3.5 and Remark 3.2]. *Assume that $\mathcal{O}$ is a connected set of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$ and $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{O}$. Let λ_* be an eigenvalue of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ with multiplicity m. Fix a small $\epsilon > 0$ such that λ_* is the only eigenvalue of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in the interval $[\lambda_* - \epsilon, \lambda_* + \epsilon]$. Then there is a connected neighborhood $\mathcal{F} \subset \mathcal{O}$ and continuous functions $\Lambda_i : \mathcal{F} \to \mathbb{R}$, $1 \leq i \leq m$, such that*

(i) *$\Lambda_i(\boldsymbol{\omega}_0, \mathbf{A}_0) = \lambda_*, 1 \leq i \leq m$;*

(ii) *$\lambda_* - \epsilon < \Lambda_1(\boldsymbol{\omega}, \mathbf{A}) \leq \cdots \leq \Lambda_m(\boldsymbol{\omega}, \mathbf{A}) < \lambda_* + \epsilon$ for each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{F}$;*

(iii) *$\Lambda_i(\boldsymbol{\omega}, \mathbf{A}), 1 \leq i \leq m$, are eigenvalues of $(\boldsymbol{\omega}, \mathbf{A})$ for each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{F}$.*

Remark 2.1. In Lemma 2.6, $\Lambda_i, 1 \leq i \leq m$, are called the continuous eigenvalue branches through λ_*.

2.3. Properties of the eigenvalue functions

In this subsection, a necessary and sufficient condition for all the eigenvalue functions to be continuous and several other properties of the eigenvalue functions in a set of the space of the SLPs, which are useful in the study of asymptotic behaviors of the n-th eigenvalue function near a discontinuity point, are obtained.

Theorem 2.1. *Let $\mathcal{O}$ be a set of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$. Then the number of eigenvalues of each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ is equal if and only if each eigenvalue function $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ restricted in $\mathcal{O}$ is continuous. Furthermore, if $\mathcal{O}$ is a connected set of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$, then each eigenvalue function $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ is locally a continuous eigenvalue branch in $\mathcal{O}$.*

Proof. Suppose that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ has exactly k eigenvalues: $\lambda_0(\boldsymbol{\omega}, \mathbf{A}) \leq \lambda_1(\boldsymbol{\omega}, \mathbf{A}) \leq \cdots \leq \lambda_{k-1}(\boldsymbol{\omega}, \mathbf{A})$. Then the k eigenvalue functions $\lambda_i : \mathcal{O} \to \mathbb{R}, 0 \leq i \leq k-1$, are well-defined. Now, we show that λ_i restricted in $\mathcal{O}$ is continuous at a fixed $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{O}$. We first consider the case that $\lambda_i(\boldsymbol{\omega}_0, \mathbf{A}_0)$ is a simple eigenvalue. Fix $0 < i < k-1$. Let $r_j, 1 \leq j \leq 4$, be four real numbers such that $r_1 < \lambda_0(\boldsymbol{\omega}_0, \mathbf{A}_0)$, $\lambda_{i-1}(\boldsymbol{\omega}_0, \mathbf{A}_0) < r_2 < \lambda_i(\boldsymbol{\omega}_0, \mathbf{A}_0) < r_3 < \lambda_{i+1}(\boldsymbol{\omega}_0, \mathbf{A}_0)$ and $r_4 > \lambda_{k-1}(\boldsymbol{\omega}_0, \mathbf{A}_0)$. By Lemma 2.5 there exists a neighborhood $\mathcal{U}$ of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in $\mathcal{O}$ such that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{U}$ has exactly i eigenvalues in (r_1, r_2), exactly one eigenvalue in (r_2, r_3), and exactly $k-i-1$ eigenvalues in (r_3, r_4). Since each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{U}$ has exactly k eigenvalues, the eigenvalue of $(\boldsymbol{\omega}, \mathbf{A})$ in (r_2, r_3) is exactly $\lambda_i(\boldsymbol{\omega}, \mathbf{A})$. Hence, λ_i restricted in $\mathcal{O}$ is continuous at $(\boldsymbol{\omega}_0, \mathbf{A}_0)$. With a similar method used above, one can easily verify that λ_0 and λ_{k-1} restricted in $\mathcal{O}$ are continuous at $(\boldsymbol{\omega}_0, \mathbf{A}_0)$.

Suppose that $\mathcal{O}$ is connected. Then the above $\mathcal{U}$ can be chosen to be also connected. By Lemma 2.6 and Remark 2.1, λ_i restricted in $\mathcal{U}$ is exactly a continuous eigenvalue branch through $\lambda_i(\boldsymbol{\omega}_0, \mathbf{A}_0)$. Thus, λ_i is locally a continuous eigenvalue branch in $\mathcal{O}$ for each $0 \leq i \leq k-1$.

In the case that the multiplicity of $\lambda_i(\boldsymbol{\omega}_0, \mathbf{A}_0)$ is equal to 2, one can show that the results still hold with a similar argument.

Conversely, suppose that there exists an $(\boldsymbol{\omega}_1, \mathbf{A}_1) \in \mathcal{O}$ such that the number of eigenvalues of $(\boldsymbol{\omega}_1, \mathbf{A}_1)$ is not equal to that of another. Then there exists an eigenvalue function λ_{i_0} that can not be well-defined at least at one point in $\mathcal{O}$. Thus λ_{i_0} restricted in $\mathcal{O}$ is not continuous. This completes the proof.

Remark 2.2. By Theorem 2.1, if the number of eigenvalues of each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ is not equal, then there exists at least one eigenvalue function that is not continuous in $\mathcal{O}$. In this case, are all the eigenvalue functions not continuous in $\mathcal{O}$? The following example gives a negative answer.

Example 2.1. Let $s \in [0, 2]$. Consider the 1-parameter family of SLPs (1.1)–(1.2) with

$$f_0 = \begin{cases} \frac{1}{2-s} & \text{if } s \in [0,1), \\ \frac{1}{s} & \text{if } s \in [1,2], \end{cases} \qquad f_1 = \begin{cases} 1 & \text{if } s \in [0,1), \\ \frac{1}{s} & \text{if } s \in [1,2], \end{cases}$$

$$f_2 = 1,\ q_1 = q_2 = 0,\ w_1 = w_2 = 1,\ N = 2,$$

and

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 0 & -1 & 1 \end{bmatrix}. \tag{2.6}$$

Then, by Lemma 2.3, direct calculations deduce that the characteristic function is

$$\Gamma(\lambda) = \begin{cases} (1-s)\lambda^2 + (s-2)\lambda + s - 1 & \text{if } s \in [0,1), \\ (s^2 - s)\lambda^2 + (s^2 - 4s + 2)\lambda + 2 - 2s & \text{if } s \in [1,2]. \end{cases}$$

Thus, the SLP with $s=1$ has exactly one eigenvalue and with each $s\in[0,1)\cup(1,2]$ has exactly two eigenvalues, which are given as

$$\lambda_0(s)=\begin{cases}\frac{2-s-\sqrt{5s^2-12s+8}}{2(1-s)} & \text{if } s\in[0,1),\\ 0 & \text{if } s=1,\\ \frac{-s^2+4s-2-\sqrt{(s^2-4s+2)^2-4(s^2-s)(2-2s)}}{2(s^2-s)} & \text{if } s\in(1,2],\end{cases}$$

$$\lambda_1(s)=\begin{cases}\frac{2-s+\sqrt{5s^2-12s+8}}{2(1-s)} & \text{if } s\in[0,1),\\ \frac{-s^2+4s-2+\sqrt{(s^2-4s+2)^2-4(s^2-s)(2-2s)}}{2(s^2-s)} & \text{if } s\in(1,2].\end{cases}$$



Figure 2.1.

See Figure 2.1. Then the number of eigenvalues of the SLP with $s=1$ is not equal to that of another one. It is evident that the eigenvalue function λ_0 is continuous in $s\in[0,2]$. Note that λ_0 is bounded from below for $s\in[0,2]$. However, the eigenvalue function λ_1 is not well-defined at $s=1$, and thus λ_1 is not continuous at $s=1$.

Now we give several other properties of the eigenvalue functions, which are useful in the discussions in the sequent sections.

Theorem 2.2. *Assume that $\mathcal{O}\subset\Omega_N^{\mathbb{R},+}\times\mathcal{B}^{\mathbb{C}}$ satisfies that each $(\boldsymbol{\omega},\mathbf{A})$ in $\mathcal{O}$ has exactly k eigenvalues, where $k\geq 2$. Let $(\boldsymbol{\omega}_0,\mathbf{A}_0)\in\bar{\mathcal{O}}\backslash\mathcal{O}$ have exactly m eigenvalues for some $0<m\leq k-1$.*

(i) *If the first $k-m$ eigenvalue functions satisfy*

$$\lim_{\mathcal{O}\ni(\boldsymbol{\omega},\mathbf{A})\to(\boldsymbol{\omega}_0,\mathbf{A}_0)}\lambda_n(\boldsymbol{\omega},\mathbf{A})=-\infty,\quad 0\leq n\leq k-m-1, \tag{2.7}$$

then the last m eigenvalue functions satisfy

$$\lim_{\mathcal{O}\ni(\boldsymbol{\omega},\mathbf{A})\to(\boldsymbol{\omega}_0,\mathbf{A}_0)}\lambda_n(\boldsymbol{\omega},\mathbf{A})=\lambda_{n-k+m}(\boldsymbol{\omega}_0,\mathbf{A}_0),\quad k-m\leq n\leq k-1. \tag{2.8}$$

(ii) *If the last $k-m$ eigenvalue functions satisfy*

$$\lim_{\mathcal{O}\ni(\boldsymbol{\omega},\mathbf{A})\to(\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega},\mathbf{A}) = +\infty, \ \ m \le n \le k-1,$$

then the first m eigenvalue functions satisfy

$$\lim_{\mathcal{O}\ni(\boldsymbol{\omega},\mathbf{A})\to(\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega},\mathbf{A}) = \lambda_n(\boldsymbol{\omega}_0,\mathbf{A}_0), \ \ 0 \le n \le m-1.$$

Proof. First, we show that (i) holds. Let $r_1 < r_2$ be any two real numbers such that $r_1 < \lambda_0(\boldsymbol{\omega}_0,\mathbf{A}_0)$ and $r_2 > \lambda_{m-1}(\boldsymbol{\omega}_0,\mathbf{A}_0)$. It follows from (2.7) that there exists a neighborhood $\mathcal{U}$ of $(\boldsymbol{\omega}_0,\mathbf{A}_0)$ in $\mathcal{O}\cup\{(\boldsymbol{\omega}_0,\mathbf{A}_0)\}$ such that

$$\lambda_n(\boldsymbol{\omega},\mathbf{A}) < r_1, \ \forall\ (\boldsymbol{\omega},\mathbf{A}) \in \mathcal{U}\backslash\{(\boldsymbol{\omega}_0,\mathbf{A}_0)\}, \ 0 \le n \le k-m-1. \tag{2.9}$$

Since $(\boldsymbol{\omega}_0,\mathbf{A}_0)$ has exactly m eigenvalues, by Lemma 2.5 there exists a neighborhood $\mathcal{U}_1 \subset \mathcal{U}$ of $(\boldsymbol{\omega}_0,\mathbf{A}_0)$ such that each $(\boldsymbol{\omega},\mathbf{A}) \in \mathcal{U}_1$ has exactly m eigenvalues in $[r_1,r_2]$, which are all in (r_1,r_2). Hence, for each $(\boldsymbol{\omega},\mathbf{A}) \in \mathcal{U}_1\backslash\{(\boldsymbol{\omega}_0,\mathbf{A}_0)\}$, by (2.9) and noting that $(\boldsymbol{\omega},\mathbf{A})$ has exactly k eigenvalues, one has that

$$\lambda_j(\boldsymbol{\omega},\mathbf{A}) \in (r_1,r_2), \ \ k-m \le j \le k-1. \tag{2.10}$$

Now, suppose that $\lambda_i(\boldsymbol{\omega}_0,\mathbf{A}_0)$ is a simple eigenvalue for some $0 \le i \le m-1$. Let $(r_3,r_4) \subset (r_1,r_2)$ be an interval such that $\lambda_i(\boldsymbol{\omega}_0,\mathbf{A}_0)$ is the only eigenvalue of $(\boldsymbol{\omega}_0,\mathbf{A}_0)$ in $[r_3,r_4]$, and $\lambda_i(\boldsymbol{\omega}_0,\mathbf{A}_0) \in (r_3,r_4)$. Again, by Lemma 2.5 there exists a neighborhood $\mathcal{U}_2 \subset \mathcal{U}_1$ of $(\boldsymbol{\omega}_0,\mathbf{A}_0)$ such that each $(\boldsymbol{\omega},\mathbf{A}) \in \mathcal{U}_2$ has exactly i eigenvalues in (r_1,r_3), exactly one eigenvalue in (r_3,r_4), and exactly $m-i-1$ eigenvalues in (r_4,r_2). Thus, it follows from (2.9) and (2.10) that $\lambda_{k-m+i}(\boldsymbol{\omega},\mathbf{A}) \in (r_3,r_4)$ for each $(\boldsymbol{\omega},\mathbf{A}) \in \mathcal{U}_2\backslash\{(\boldsymbol{\omega}_0,\mathbf{A}_0)\}$. Consequently,

$$\lim_{\mathcal{O}\ni(\boldsymbol{\omega},\mathbf{A})\to(\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_{k-m+i}(\boldsymbol{\omega},\mathbf{A}) = \lambda_i(\boldsymbol{\omega}_0,\mathbf{A}_0).$$

Therefore, (2.8) holds in this case.

In the other case that the multiplicity of $\lambda_i(\boldsymbol{\omega}_0,\mathbf{A}_0)$ is equal to 2 for some $0 \le i \le m-1$, one can show that the results hold with a similar argument.

Since the proof of (ii) is similar to that of (i), we omit its details. The proof is complete.

Theorem 2.3. *Let $\mathcal{O}$ be a connected set of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$ and r_1 and r_2 be two real numbers with $r_1 < r_2$. Assume that each problem in $\mathcal{O}$ has exactly k eigenvalues, and exactly $m \ge 0$ eigenvalues in $[r_1,r_2]$ that are in (r_1,r_2) with $m < k$. Then its other $k-m$ eigenvalues out of $[r_1,r_2]$, denoted by $\hat\lambda_1(\boldsymbol{\omega},\mathbf{A}) \le \cdots \le \hat\lambda_{k-m}(\boldsymbol{\omega},\mathbf{A})$, have the following properties:*

(i) *For each $1 \le i \le k-m$,*

$$\textit{either } E_i := \{\hat\lambda_i(\boldsymbol{\omega},\mathbf{A}) : (\boldsymbol{\omega},\mathbf{A}) \in \mathcal{O}\} \subset (-\infty,r_1) \textit{ or } E_i \subset (r_2,+\infty); \tag{2.11}$$

there exists $0 \le i_0 \le k-1$ such that $\hat\lambda_i(\boldsymbol{\omega},\mathbf{A}) = \lambda_{i_0}(\boldsymbol{\omega},\mathbf{A})$ for all $(\boldsymbol{\omega},\mathbf{A}) \in \mathcal{O}$, and consequently, $\hat\lambda_i$ is continuous in $\mathcal{O}$.

(ii) *Further, assume that* $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \bar{\mathcal{O}} \backslash \mathcal{O}$ *has exactly* m *eigenvalues* $\lambda_j(\boldsymbol{\omega}_0, \mathbf{A}_0)$*,* $0 \le j \le m-1$*, and all of them are in* (r_1, r_2)*. If* $E_i \subset (-\infty, r_1)$ *for some* $1 \le i \le k-m$*, then*

$$\lim_{\mathcal{O} \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \hat{\lambda}_i(\boldsymbol{\omega}, \mathbf{A}) = -\infty.$$

If $E_i \subset (r_2, +\infty)$ *for some* $1 \le i \le k-m$*, then*

$$\lim_{\mathcal{O} \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \hat{\lambda}_i(\boldsymbol{\omega}, \mathbf{A}) = +\infty.$$

Proof. By the assumption that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ has exactly k eigenvalues, the n-th eigenvalue functions λ_j, $0 \le j \le k-1$, restricted in $\mathcal{O}$ are continuous by Theorem 2.1.

First, we show that (i) holds. We claim that (2.11) holds for $i = k-m$. Otherwise, there exist $(\boldsymbol{\omega}_1, \mathbf{A}_1)$, $(\boldsymbol{\omega}_2, \mathbf{A}_2) \in \mathcal{O}$ such that $\hat{\lambda}_{k-m}(\boldsymbol{\omega}_1, \mathbf{A}_1) \in (-\infty, r_1)$ and $\hat{\lambda}_{k-m}(\boldsymbol{\omega}_2, \mathbf{A}_2) \in (r_2, +\infty)$. Then $\lambda_{k-1}(\boldsymbol{\omega}_1, \mathbf{A}_1) \in (r_1, r_2)$ and $\lambda_{k-1}(\boldsymbol{\omega}_2, \mathbf{A}_2) \in (r_2, +\infty)$. Since r_2 is not an eigenvalue of any problem in $\mathcal{O}$, λ_{k-1} is not continuous in $\mathcal{O}$. This is a contradiction by the assumption that $\mathcal{O}$ is connected.

If $E_{k-m} \subset (-\infty, r_1)$, then $E_i \subset (-\infty, r_1)$ for all $1 \le i \le k-m-1$, and consequently, $\hat{\lambda}_j(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{j-1}(\boldsymbol{\omega}, \mathbf{A})$ for all $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ and $1 \le j \le k-m$.

If $E_{k-m} \subset (r_2, +\infty)$, then using the same method as employed in the above paragraph, one can show that either $E_{k-m-1} \subset (-\infty, r_1)$ or $E_{k-m-1} \subset (r_2, +\infty)$. If $E_{k-m-1} \subset (-\infty, r_1)$, then $E_j \subset (-\infty, r_1)$, $1 \le j \le k-m-2$. Hence, $\hat{\lambda}_{k-m}(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{k-1}(\boldsymbol{\omega}, \mathbf{A})$ and $\hat{\lambda}_j(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{j-1}(\boldsymbol{\omega}, \mathbf{A})$, $1 \le j \le k-m-1$, for all $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$. If $E_{k-m-1} \subset (r_2, +\infty)$, then again using the same method as employed in the above discussion, one can show that either $E_{k-m-2} \subset (-\infty, r_1)$ or $E_{k-m-2} \subset (r_2, +\infty)$. This procedure can be finished in finite steps.

Further, since the n-th eigenvalue functions $\lambda_j, 0 \le j \le k-1$, restricted in $\mathcal{O}$ are continuous, $\hat{\lambda}_i$ is continuous in $\mathcal{O}$ for each $1 \le i \le k-m$. Then (i) has been shown.

Now, we show that (ii) holds. We only consider the case that $E_i \subset (-\infty, r_1)$ for some $1 \le i \le k-m$. The other case can be similarly discussed. Otherwise, there exists a positive number $M > |r_1|$ such that for any neighborhood $\mathcal{U}$ of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in $\mathcal{O} \cup \{(\boldsymbol{\omega}_0, \mathbf{A}_0)\}$, there exists $(\hat{\boldsymbol{\omega}}, \hat{\mathbf{A}}) \in \mathcal{U} \backslash \{(\boldsymbol{\omega}_0, \mathbf{A}_0)\}$ satisfying $-M \le \hat{\lambda}_i(\hat{\boldsymbol{\omega}}, \hat{\mathbf{A}}) < r_1$. Since $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ has exactly m eigenvalues in $(-M, r_2)$, by Lemma 2.5 there exists a neighborhood $\mathcal{U}_1 \subset \mathcal{O} \cup \{(\boldsymbol{\omega}_0, \mathbf{A}_0)\}$ of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ such that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{U}_1$ has exactly m eigenvalues in $(-M, r_2)$. However, taking $\mathcal{U} \subset \mathcal{U}_1$, we get that $(\hat{\boldsymbol{\omega}}, \hat{\mathbf{A}})$ has at least $m+1$ eigenvalues in $(-M, r_2)$ by the assumption that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{O}$ has m eigenvalues in (r_1, r_2). This is a contradiction. The entire proof is complete.

In some cases, $(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$ may be continuously dependent on some real parameters or variables. For example, $(\boldsymbol{\omega}, \mathbf{A})$ is continuously dependent on $1/f_0$; $(\boldsymbol{\omega}, \mathbf{S}_{\alpha,\beta})$

is continuously dependent on α and β; $(\boldsymbol{\omega}, \mathbf{A})$, where $\mathbf{A} \in \mathcal{O}^{\mathbb{C}}_{1,3}$, is continuously dependent on variables a_{12} and b_{22}, etc. Then, we write $(\boldsymbol{\omega}, \mathbf{A})_\nu$ instead of $(\boldsymbol{\omega}, \mathbf{A})$ to indicate the dependence of $(\boldsymbol{\omega}, \mathbf{A})$ on a variable or parameter ν in some situations for convenience. Next, we shall discuss the dependence of $\lambda_n(\nu) := \lambda_n((\boldsymbol{\omega}, \mathbf{A})_\nu)$ on ν.

Lemma 2.7. *Let $(\boldsymbol{\omega}, \mathbf{A})_\nu \in \Omega^{\mathbb{R},+}_N \times \mathcal{B}^{\mathbb{C}}$ be continuously dependent on a real variable or parameter ν in $(\nu_0 - \epsilon, \nu_0 + \epsilon)$ for some $\epsilon > 0$ and $\mathcal{O} := \{(\boldsymbol{\omega}, \mathbf{A})_\nu : \nu \in (\nu_0 - \epsilon, \nu_0 + \epsilon)\}$. Assume that $(\boldsymbol{\omega}, \mathbf{A})_{\nu_0}$ has exactly $m \ge 1$ eigenvalues and for each $\nu \in (\nu_0 - \epsilon, \nu_0 + \epsilon)\backslash\{\nu_0\}$, $(\boldsymbol{\omega}, \mathbf{A})_\nu$ has exactly k eigenvalues with $k > m$.*

(i) *If the n-th eigenvalue functions $\lambda_n(\nu)$ are non-increasing in $(\nu_0 - \epsilon_0, \nu_0)$ for all $0 \le n \le k-1$, then they satisfy the following asymptotic behaviors near ν_0:*

$$\lim_{\nu\to\nu_0^-} \lambda_n(\nu) = -\infty, \ 0 \le n \le k-m-1, \tag{2.12}$$

$$\lim_{\nu\to\nu_0^-} \lambda_n(\nu) = \lambda_{n-k+m}(\nu_0), \ k-m \le n \le k-1. \tag{2.13}$$

(ii) *If the n-th eigenvalue functions $\lambda_n(\nu)$ are non-decreasing in $(\nu_0 - \epsilon_0, \nu_0)$ for all $0 \le n \le k-1$, then they satisfy the following asymptotic behaviors near ν_0:*

$$\lim_{\nu\to\nu_0^-} \lambda_n(\nu) = \lambda_n(\nu_0), \ \ 0 \le n \le m-1, \ \lim_{\nu\to\nu_0^-} \lambda_n(\nu) = +\infty, \ m \le n \le k-1.$$

(iii) *If the n-th eigenvalue functions $\lambda_n(\nu)$ are non-increasing in $(\nu_0, \nu_0 + \epsilon_0)$ for all $0 \le n \le k-1$, then they satisfy the following asymptotic behaviors near ν_0:*

$$\lim_{\nu\to\nu_0^+} \lambda_n(\nu) = \lambda_n(\nu_0), \ \ 0 \le n \le m-1, \ \lim_{\nu\to\nu_0^+} \lambda_n(\nu) = +\infty, \ m \le n \le k-1.$$

(iv) *If the n-th eigenvalue functions $\lambda_n(\nu)$ are non-decreasing in $(\nu_0, \nu_0 + \epsilon_0)$ for all $0 \le n \le k-1$, then they satisfy the following asymptotic behaviors near ν_0:*

$$\lim_{\nu\to\nu_0^+} \lambda_n(\nu) = -\infty, \ \ 0 \le n \le k-m-1, \ \lim_{\nu\to\nu_0^+} \lambda_n(\nu) = \lambda_{n-k+m}(\nu_0), \ \ k-m \le n \le k-1.$$

Proof. We only show that (i) holds. The other claims can be shown similarly.

Let (r_1, r_2) be a finite interval such that $\lambda_j(\nu_0) \in (r_1, r_2)$ for $0 \le j \le m-1$. Then, by Lemma 2.5 there exists $0 < \epsilon_0 < \epsilon$ such that for each $\nu \in (\nu_0 - \epsilon_0, \nu_0)$, $(\boldsymbol{\omega}, \mathbf{A})_\nu$ has exactly m eigenvalues in $[r_1, r_2]$, which are all in (r_1, r_2). Its other $k-m$ eigenvalues out of $[r_1, r_2]$ are denoted by $\hat{\lambda}_1(\nu) \le \cdots \le \hat{\lambda}_{k-m}(\nu)$.

Fix $1 \le i \le k-m$. We claim that $\mathcal{O}^- := \{\hat{\lambda}_i(\nu) : \nu \in (\nu_0 - \epsilon_0, \nu_0)\} \subset (-\infty, r_1)$. Otherwise, by (i) of Theorem 2.3, $\mathcal{O}^- \subset (r_2, +\infty)$ and there exists $0 \le i_0 \le k-1$ such that $\hat{\lambda}_i(\nu) = \lambda_{i_0}(\nu)$ for each $\nu \in (\nu_0 - \epsilon_0, \nu_0)$. Since $\lambda_{i_0}(\nu)$ is non-increasing in $(\nu_0 - \epsilon_0, \nu_0)$, $\hat{\lambda}_i(\nu) = \lambda_{i_0}(\nu) \le \lambda_{i_0}(\nu_0 - \epsilon_0/2) = \hat{\lambda}_i(\nu_0 - \epsilon_0/2)$ for each $\nu \in (\nu_0 - \epsilon_0/2, \nu_0)$. By Lemma 2.5, there exists $0 < \epsilon_1 < \epsilon_0/2$ such that $(\boldsymbol{\omega}, \mathbf{A})_\nu$ has exactly m eigenvalues in $(r_1, \hat{\lambda}_i(\nu_0 - \epsilon_0/2) + 1)$ for each $\nu \in (\nu_0 - \epsilon_1, \nu_0]$. Since $r_2 < \hat{\lambda}_i(\nu) < \hat{\lambda}_i(\nu_0 - \epsilon_0/2) + 1$

and $(\boldsymbol{\omega}, \mathbf{A})_\nu$ has exactly m eigenvalues in (r_1, r_2) for each $\nu \in (\nu_0 - \epsilon_1, \nu_0)$, it follows that $(\boldsymbol{\omega}, \mathbf{A})_\nu$ has at least $m+1$ eigenvalues in $(r_1, \hat{\lambda}_i(\nu_0 - \epsilon_0/2) + 1)$ for each $\nu \in (\nu_0 - \epsilon_1, \nu_0)$. This is a contradiction. Hence the claim holds.

Consequently, $\hat{\lambda}_n(\nu) = \lambda_{n-1}(\nu)$, $\nu \in (\nu_0 - \epsilon_0, \nu_0)$, for each $1 \le n \le k - m$ again by (i) of Theorem 2.3. It follows from (ii) of Theorem 2.3 that (2.12) holds and then (2.13) holds by (i) of Theorem 2.2. The proof is complete.

Remark 2.3. If $m = 0$ and the conditions in (i) of Lemma 2.7 are satisfied, then the n-th eigenvalue functions $\lambda_n(\nu)$, $0 \le n \le k-1$, satisfy the following asymptotic behaviors near ν_0:

$$\lim_{\nu \to \nu_0^-} \lambda_n(\nu) = -\infty, \ 0 \le n \le k-1.$$

(ii)–(iv) in Lemma 2.7 can be modified similarly in the case that $m = 0$.

Remark 2.4. If the conditions in (i) and (iii) (or (ii) and (iv)) of Lemma 2.7 are satisfied, then $(\boldsymbol{\omega}, \mathbf{A})_{\nu_0}$ is a discontinuity point of λ_n for all $0 \le n \le k-1$.

Remark 2.5. In Example 2.1, $\lambda_n(s)$ is non-decreasing in $s \in [0, 1)$, and non-increasing in $s \in (1, 2]$ for each $n = 0, 1$. Thus, the conditions in (ii)–(iii) of Lemma 2.7 are satisfied for $k = 2$ and $m = 1$, and therefore,

$$\lim_{s \to 1^-} \lambda_0(s) = \lim_{s \to 1^+} \lambda_0(s) = \lambda_0(1), \ \lim_{s \to 1^-} \lambda_1(s) = +\infty, \ \lim_{s \to 1^+} \lambda_1(s) = +\infty.$$

This also shows that $\lambda_0(s)$ is continuous at $s = 1$, and $\lambda_1(s)$ is not continuous at $s = 1$.

3. Continuity and discontinuity of the n-th eigenvalue function in the space of the SLEs

In this section, the continuous and discontinuous dependence of the n-th eigenvalue function on equation (1.1) is discussed. Its continuity and discontinuity sets in $\Omega_N^{\mathbb{R},+}$ are given and its monotonicity in some directions in the continuity set is studied. In particular, its asymptotic behaviors near a discontinuity point are completely characterized.

It was shown in [10] that the n-th eigenvalue depends continuously on the differential equation (1.5) in the continuous case. However, the following example shows that the n-th eigenvalue may not depend continuously on equation (1.1) in the discrete case.

Example 3.1. Let $s \in [1/10, 2]$. Consider the 1-parameter family of the SLPs, in which

$$f_0 = 1/s, \ f_1 = f_2 = 1, \ q_1 = q_2 = 0, \ w_1 = w_2 = 1, \ N = 2,$$

and the coefficients in (1.2) is the same as (2.6). Then, by Lemma 2.3, direct calculations deduce that the characteristic function is

$$\Gamma(\lambda) = (s-1)\lambda^2 - s\lambda + 1 - s.$$

Thus, the SLP with $s = 1$ has exactly one eigenvalue and with each $s \in [1/10, 1) \cup (1, 2]$ has exactly two eigenvalues, which are given as

$$\lambda_0(s) = \begin{cases} \frac{s+\sqrt{5s^2-8s+4}}{2(s-1)} & \text{if } s \in [1/10, 1), \\ 0 & \text{if } s = 1, \\ \frac{s-\sqrt{5s^2-8s+4}}{2(s-1)} & \text{if } s \in (1, 2], \end{cases} \qquad \lambda_1(s) = \begin{cases} \frac{s-\sqrt{5s^2-8s+4}}{2(s-1)} & \text{if } s \in [1/10, 1), \\ \frac{s+\sqrt{5s^2-8s+4)}}{2(s-1)} & \text{if } s \in (1, 2]. \end{cases}$$

See Figure 3.1 below. It is evident that the n-th eigenvalue function λ_n is not continuous at $s = 1$ for each $n = 0, 1$. This example shows that the n-th eigenvalue may not depend continuously on $1/f_0$ in general.



Figure 3.1.

Fix a BC

$$\hat{\mathbf{A}} = \begin{bmatrix} \hat{a}_{11} & \hat{a}_{12} & \hat{b}_{11} & \hat{b}_{12} \\ \hat{a}_{21} & \hat{a}_{22} & \hat{b}_{21} & \hat{b}_{22} \end{bmatrix}$$

in this section. We shall study continuous dependence of the n-th eigenvalue on the SLE.

We need the following monotonicity results of the continuous eigenvalue branches:

Lemma 3.1 [22, Theorem 4.8]. *Fix a BC* $\hat{\mathbf{A}}$*. Then, each continuous eigenvalue branch* Λ *over* $\Omega_N^{\mathbb{R},+}$ *is non-increasing in every* $(1/f_j)$*-direction for* $0 \le j \le N-1$*, independent of* f_N*, and non-decreasing in every* q_j*-direction; while its positive and negative parts are non-increasing and non-decreasing in every* w_j*-direction, respectively.*

For convenience, we introduce the following notations. Let $\mu_1 := \hat{a}_{11}\hat{b}_{22} - \hat{a}_{21}\hat{b}_{12}$, $\mu_2 := \hat{a}_{22}\hat{b}_{12} - \hat{a}_{12}\hat{b}_{22}$. If $\mu_1 \neq 0$, denote

$$\eta := -\mu_2/\mu_1, \ \mathcal{E} := \{\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} : 1/f_0 = \eta\},$$
$$\mathcal{E}^+ := \{\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} : 1/f_0 > \eta\}, \ \mathcal{E}^- := \{\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} : 1/f_0 < \eta\},$$
$$\mathcal{E}_1 := \{\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} : 1/f_0 = -1/\hat{a}_{11}\} \text{ if } \hat{a}_{11} \neq 0; \ \ \mathcal{E}_2 := \{\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} : 1/f_0 = \hat{a}_{12}\}.$$

And $\mathcal{E}_1^+$, $\mathcal{E}_1^-$, $\mathcal{E}_2^+$, $\mathcal{E}_2^-$ can be similarly defined.

Due to (2.5) and the discussion below it, we shall divide our study here into the following three cases:

(i) $\mu_1 \neq 0, \mu_2 \neq 0$;

(ii) either $\mu_1 = 0, \mu_2 \neq 0$, or $\mu_1 \neq 0, \mu_2 = 0$;

(iii) $\mu_1 = 0, \mu_2 = 0$.

Theorem 3.1. *Fix a BC* $\hat{\mathbf{A}}$. *Assume that* $\mu_1 \neq 0$ *and* $\mu_2 \neq 0$. *Then for each* $\boldsymbol{\omega} \in \mathcal{E}$, $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ *has exactly* $N-1$ *eigenvalues, and for each* $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$, $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ *has exactly* N *eigenvalues* $\lambda_n(\boldsymbol{\omega}) := \lambda_n(\boldsymbol{\omega}, \hat{\mathbf{A}})$, $0 \leq n \leq N-1$, *which satisfy that*

(i) $\lambda_n(\boldsymbol{\omega})$ *are continuous in* $\Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$*;*

(ii) $\lambda_n(\boldsymbol{\omega})$ *restricted in each connected component of* $\mathcal{E}^+$ *or* $\mathcal{E}^-$ *are non-increasing in every* $(1/f_j)$*-direction, independent of* f_N*, and non-decreasing in every* q_j*-direction, while its positive and negative parts are non-increasing and non-decreasing in every* w_j*-direction, respectively;*

(iii) $\lambda_n(\boldsymbol{\omega})$ *are not continuous at each point of* $\mathcal{E}$*, and have the following asymptotic behaviors near any given* $\boldsymbol{\omega}_0 \in \mathcal{E}$*:*

$$\lim_{\mathcal{E}^- \ni \boldsymbol{\omega} \to \boldsymbol{\omega}_0} \lambda_0(\boldsymbol{\omega}) = -\infty, \tag{3.1}$$

$$\lim_{\mathcal{E}^- \ni \boldsymbol{\omega} \to \boldsymbol{\omega}_0} \lambda_n(\boldsymbol{\omega}) = \lambda_{n-1}(\boldsymbol{\omega}_0), \quad 1 \leq n \leq N-1, \tag{3.2}$$

$$\lim_{\mathcal{E}^+ \cup \mathcal{E} \ni \boldsymbol{\omega} \to \boldsymbol{\omega}_0} \lambda_n(\boldsymbol{\omega}) = \lambda_n(\boldsymbol{\omega}_0), \quad 0 \leq n \leq N-2, \tag{3.3}$$

$$\lim_{\mathcal{E}^+ \ni \boldsymbol{\omega} \to \boldsymbol{\omega}_0} \lambda_{N-1}(\boldsymbol{\omega}) = +\infty. \tag{3.4}$$

And consequently, $\lambda_n(\boldsymbol{\omega})$ *restricted in* $\mathcal{E}^+ \cup \mathcal{E}$ *is continuous for each* $0 \leq n \leq N-2$.

Proof. By (2.4), $r(\boldsymbol{\omega}, \hat{\mathbf{A}}) = 1$ for each $\boldsymbol{\omega} \in \mathcal{E}$, and $r(\boldsymbol{\omega}, \hat{\mathbf{A}}) = 2$ for each $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$. It follows from Lemma 2.4 that $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly $N-1$ eigenvalues for each $\boldsymbol{\omega} \in \mathcal{E}$ and $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly N eigenvalues for each $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$.

Now, we show that (i) and (ii) hold. Since $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly N eigenvalues $\lambda_n(\boldsymbol{\omega})$, $0 \leq n \leq N-1$, for each $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$, which is an open set of $\Omega_N^{\mathbb{R},+}$, $\lambda_n(\boldsymbol{\omega})$ is continuous in $\Omega_N^{\mathbb{R},+} \backslash \mathcal{E}$ and is locally a continuous eigenvalue branch in each connected component of $\mathcal{E}^+$ or $\mathcal{E}^-$ for each $0 \leq n \leq N-1$ by Theorem 2.1. This, together with Lemma 3.1, implies its monotonicity in each connected component of $\mathcal{E}^+$ or $\mathcal{E}^-$.

Next, we show that (iii) holds. It suffices to prove (3.1)–(3.4). Fix any

$$\boldsymbol{\omega}_0 := (\eta, 1/f_0^0, \cdots, 1/f_N^0, q_1^0, \cdots, q_N^0, w_1^0, \cdots, w_N^0) \in \mathcal{E}.$$

Let (r_1, r_2) be a finite interval such that $\lambda_j(\boldsymbol{\omega}_0) \in (r_1, r_2)$, $0 \leq j \leq N-2$. By Lemma 2.5, there exists a neighborhood $\mathcal{U}$ of $\boldsymbol{\omega}_0$ in $\Omega_N^{\mathbb{R},+}$ such that for each $\boldsymbol{\omega} \in \mathcal{U}$, $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly $N-1$ eigenvalues in $[r_1, r_2]$, which are all in (r_1, r_2). Denote

$$\mathcal{U}^- := \mathcal{E}^- \cap \mathcal{U},\ \mathcal{U}^0 := \mathcal{E} \cap \mathcal{U},\ \mathcal{U}^+ := \mathcal{E}^+ \cap \mathcal{U}.$$

Note that $\mathscr{U}$ can be chosen sufficiently small such that $\mathscr{U}^-$, $\mathscr{U}^0$ and $\mathscr{U}^+$ are connected. Since $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly N eigenvalues for each $\boldsymbol{\omega} \in \mathscr{U}^-$, it has exactly one eigenvalue, denoted by $\hat{\lambda}(\boldsymbol{\omega})$, outside $[r_1, r_2]$. It follows from (i) of Theorem 2.3 that either $F := \{\hat{\lambda}(\boldsymbol{\omega}) : \boldsymbol{\omega} \in \mathscr{U}^-\} \subset (-\infty, r_1)$ or $F \subset (r_2, +\infty)$.

Denote $\boldsymbol{\omega}(s) := (s, 1/f_1^0, \cdots, 1/f_N^0, q_1^0, \cdots, q_N^0, w_1^0, \cdots, w_N^0)$, $s \in \mathbb{R}$, and $\mathcal{O} := \{\boldsymbol{\omega}(s) : s \in (\eta - \epsilon, \eta)\}$ with $0 < \epsilon < |\eta|$. Then $\boldsymbol{\omega}_0 = \boldsymbol{\omega}(\eta)$ and for each $s \in (\eta - \epsilon, \eta)$, $(\boldsymbol{\omega}(s), \hat{\mathbf{A}})$ has exactly N eigenvalues $\lambda_n(\boldsymbol{\omega}(s))$, $0 \le n \le N-1$, and $(\boldsymbol{\omega}(\eta), \hat{\mathbf{A}})$ has exactly $N-1$ eigenvalues. By (ii), $\lambda_n(\boldsymbol{\omega}(s))$ is non-increasing in $(\eta - \epsilon, \eta)$ for each $0 \le n \le N-1$. Hence, $\lim_{s \to \eta^-} \lambda_0(\boldsymbol{\omega}(s)) = -\infty$ by (i) of Lemma 2.7. This implies that there exists an $\boldsymbol{\omega}_1 \in \mathscr{U}^-$ such that $\lambda_0(\boldsymbol{\omega}_1) < r_1$. Hence, $\hat{\lambda}(\boldsymbol{\omega}_1) = \lambda_0(\boldsymbol{\omega}_1)$. Thus, again by (i) of Theorem 2.3, $F = \{\lambda_0(\boldsymbol{\omega}) : \boldsymbol{\omega} \in \mathscr{U}^-\} \subset (-\infty, r_1)$. By (ii) of Theorem 2.3 one gets that (3.1) holds.

With a similar argument to the proof of (3.1), one can show that (3.4) holds.

Note that $\mathscr{U}^0$ is connected and $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ has exactly $N-1$ eigenvalues for each $\boldsymbol{\omega} \in \mathscr{U}^0$. Hence, λ_n restricted in $\mathscr{U}^0$ is continuous and locally a continuous eigenvalue branch for each $0 \le n \le N-2$ by Theorem 2.1. This, together with Theorem 2.2, implies that (3.2) and (3.3) hold. The entire proof is complete.

Theorem 3.2. *Fix a BC* $\hat{\mathbf{A}}$. *Assume that either* $\mu_1 = 0, \mu_2 \neq 0$ *or* $\mu_1 \neq 0, \mu_2 = 0$. *Then for each* $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+}$, $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ *has exactly* N *eigenvalues* $\lambda_n(\boldsymbol{\omega})$, $0 \le n \le N-1$, *which satisfy that*

- (i) $\lambda_n(\boldsymbol{\omega})$ *are continuous in* $\Omega_N^{\mathbb{R},+}$;
- (ii) $\lambda_n(\boldsymbol{\omega})$ *restricted in each connected component of* $\Omega_N^{\mathbb{R},+}$ *have the same monotonicity as that in* (ii) *of Theorem* 3.1.

Theorem 3.3. *Fix a BC* $\hat{\mathbf{A}}$. *Assume that* $\mu_1 = 0$ *and* $\mu_2 = 0$. *Then the BC* $\hat{\mathbf{A}}$ *can be written as*

$$\text{either} \quad \hat{\mathbf{A}}_1 = \begin{bmatrix} \hat{a}_{11} & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix} \quad \text{or} \quad \hat{\mathbf{A}}_2 = \begin{bmatrix} 1 & \hat{a}_{12} & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix}.$$

- (i) *In the case that* $\hat{\mathbf{A}} = \hat{\mathbf{A}}_1$ *with* $\hat{a}_{11} \neq 0$, $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ *has exactly* $N-2$ *eigenvalues for each* $\boldsymbol{\omega} \in \mathscr{E}_1$, *and* $(\boldsymbol{\omega}, \hat{\mathbf{A}})$ *has exactly* $N-1$ *eigenvalues* $\lambda_n(\boldsymbol{\omega})$, $0 \le n \le N-2$, *for each* $\boldsymbol{\omega} \in \Omega_N^{\mathbb{R},+} \backslash \mathscr{E}_1$, *which satisfy that*
  - (ia) $\lambda_n(\boldsymbol{\omega})$ *are continuous in* $\Omega_N^{\mathbb{R},+} \backslash \mathscr{E}_1$;
  - (ib) $\lambda_n(\boldsymbol{\omega})$ *restricted in each connected component of* $\mathscr{E}_1^+$ *or* $\mathscr{E}_1^-$ *have the same monotonicity as that in* (ii) *of Theorem* 3.1;
  - (ic) $\lambda_n(\boldsymbol{\omega})$ *are not continuous at each point of* $\mathscr{E}_1$, *and have the following asymptotic behaviors near any given* $\boldsymbol{\omega}_0 \in \mathscr{E}_1$:

$$\lim_{\mathcal{E}_1^-\ni\boldsymbol{\omega}\to\boldsymbol{\omega}_0}\lambda_0(\boldsymbol{\omega})=-\infty,\ \lim_{\mathcal{E}_1^-\ni\boldsymbol{\omega}\to\boldsymbol{\omega}_0}\lambda_n(\boldsymbol{\omega})=\lambda_{n-1}(\boldsymbol{\omega}_0),\ 1\le n\le N-2,$$
$$\lim_{\mathcal{E}_1^+\cup\mathcal{E}_1\ni\boldsymbol{\omega}\to\boldsymbol{\omega}_0}\lambda_n(\boldsymbol{\omega})=\lambda_n(\boldsymbol{\omega}_0),\ 0\le n\le N-3,\ \lim_{\mathcal{E}_1^+\ni\boldsymbol{\omega}\to\boldsymbol{\omega}_0}\lambda_{N-2}(\boldsymbol{\omega})=+\infty.$$

And consequently, $\lambda_n(\boldsymbol{\omega})$ *restricted in* $\mathcal{E}_1^+\cup\mathcal{E}_1$ *is continuous for each* $0\le n\le N-3$.

(ii) *In the case that* $\hat{\mathbf{A}}=\hat{\mathbf{A}}_2$ *with* $\hat{a}_{12}\neq 0$, *similar results in* (i) *hold for* $\mathcal{E}_1$, $\mathcal{E}_1^+$, $\mathcal{E}_1^-$ *replaced by* $\mathcal{E}_2$, $\mathcal{E}_2^+$, $\mathcal{E}_2^-$, *respectively.*

(iii) *In the case that* $\hat{\mathbf{A}}=\hat{\mathbf{A}}_1$ *with* $\hat{a}_{11}=0$ *or* $\hat{\mathbf{A}}=\hat{\mathbf{A}}_2$ *with* $\hat{a}_{12}=0$, $(\boldsymbol{\omega},\hat{\mathbf{A}})$ *has exactly* $N-1$ *eigenvalues* $\lambda_n(\boldsymbol{\omega})$, $0\le n\le N-2$, *for each* $\boldsymbol{\omega}\in\Omega_N^{\mathbb{R},+}$, *which satisfy that*

- (iiia) $\lambda_n(\boldsymbol{\omega})$ *are continuous in* $\Omega_N^{\mathbb{R},+}$;
- (iiib) $\lambda_n(\boldsymbol{\omega})$ *restricted in each connected component of* $\Omega_N^{\mathbb{R},+}$ *have the same monotonicity as that in* (ii) *of Theorem* 3.1.

Since the proofs of Theorems 3.2–3.3 are similar to that of Theorem 3.1, we omit their details.

4. Continuity and discontinuity of the n-th eigenvalue function in the space of the BCs

In this section, the continuous and discontinuous dependence of the n-th eigenvalue function on the boundary condition (1.2) is investigated. Its continuity and discontinuity sets in $\mathcal{B}^{\mathbb{C}}$ are given and its monotonicity in some directions in the continuity set is studied. Especially, its asymptotic behaviors near a discontinuity point are completely characterized.

Fix a difference equation $\hat{\boldsymbol{\omega}}=(1/\hat{f},\hat{q},\hat{w})\in\Omega_N^{\mathbb{R},+}$ in this section. By Lemma 2.1, $\mathcal{B}^{\mathbb{C}}$ equals the union of $\mathcal{O}_{1,3}^{\mathbb{C}}$, $\mathcal{O}_{2,3}^{\mathbb{C}}$, $\mathcal{O}_{1,4}^{\mathbb{C}}$ and $\mathcal{O}_{2,4}^{\mathbb{C}}$, which are four open sets of $\mathcal{B}^{\mathbb{C}}$. Thus, we shall consider the n-th eigenvalue function λ_n in $\mathcal{O}_{1,3}^{\mathbb{C}}$, $\mathcal{O}_{2,3}^{\mathbb{C}}$, $\mathcal{O}_{1,4}^{\mathbb{C}}$, and $\mathcal{O}_{2,4}^{\mathbb{C}}$, separately. We shall remark that the method used here is different from that used in the continuous case [10], where the authors divided the BCs into the separated and coupled ones. The method used here is more convenient in dealing with the discrete case. Finally, we shall apply our results to the separated and coupled BCs.

We now introduce the following notations for convenience:

$$\mathcal{B}_{1,4}:=\left\{\mathbf{A}\in\mathcal{O}_{1,4}^{\mathbb{C}}:a_{12}=1/\hat{f}_0\right\},\ \mathcal{B}_{2,4}:=\left\{\mathbf{A}\in\mathcal{O}_{2,4}^{\mathbb{C}}:a_{11}=-\hat{f}_0\right\},$$
$$\mathcal{B}_{1,3}:=\left\{\mathbf{A}\in\mathcal{O}_{1,3}^{\mathbb{C}}:(a_{12}-1/\hat{f}_0)b_{22}=|z|^2\right\},\ \mathcal{B}_{2,3}:=\left\{\mathbf{A}\in\mathcal{O}_{2,3}^{\mathbb{C}}:(a_{11}+\hat{f}_0)b_{22}=|z|^2\right\},$$
$$\mathbf{C}:=\begin{bmatrix}1&1/\hat{f}_0&0&0\\0&0&1&0\end{bmatrix},\ \mathcal{B}_{1,3r}:=\left\{\mathbf{A}\in\mathcal{B}_{1,3}:a_{12}\ge 1/\hat{f}_0,b_{22}\ge 0\right\}\setminus\{\mathbf{C}\},$$
$$\mathcal{B}_{1,3l}:=\left\{\mathbf{A}\in\mathcal{B}_{1,3}:a_{12}\le 1/\hat{f}_0,b_{22}\le 0\right\}\setminus\{\mathbf{C}\},$$

$$\begin{aligned}\mathcal{B}_{1,3r}^{+} &:= \left\{\mathbf{A} \in \mathcal{O}_{1,3}^{\mathbb{C}} : a_{12} \geq 1/\hat{f}_0, b_{22} \geq 0, (a_{12} - 1/\hat{f}_0)b_{22} > |z|^2\right\}, \\ \mathcal{B}_{1,3l}^{+} &:= \left\{\mathbf{A} \in \mathcal{O}_{1,3}^{\mathbb{C}} : a_{12} \leq 1/\hat{f}_0, b_{22} \leq 0, (a_{12} - 1/\hat{f}_0)b_{22} > |z|^2\right\}, \\ \mathcal{B}_{2,3r} &:= \left\{\mathbf{A} \in \mathcal{B}_{2,3} : a_{11} + \hat{f}_0 \geq 0, b_{22} \geq 0\right\} \setminus \{\mathbf{C}\}, \\ \mathcal{B}_{2,3l} &:= \left\{\mathbf{A} \in \mathcal{B}_{2,3} : a_{11} + \hat{f}_0 \leq 0, b_{22} \leq 0\right\} \setminus \{\mathbf{C}\},\end{aligned}$$

$\mathcal{B}_{1,4}^{+}$ and $\mathcal{B}_{2,4}^{+}$, and $\mathcal{B}_{1,4}^{-}$, $\mathcal{B}_{2,4}^{-}$, $\mathcal{B}_{1,3}^{-}$ and $\mathcal{B}_{2,3}^{-}$ can be defined similarly as $\mathcal{E}^{+}$ and $\mathcal{E}^{-}$ in Section 3, respectively; $\mathcal{B}_{2,3r}^{+}$ and $\mathcal{B}_{2,3l}^{+}$ can be defined similarly as $\mathcal{B}_{1,3r}^{+}$ and $\mathcal{B}_{1,3l}^{+}$, respectively. Then $\mathcal{O}_{i,4}^{\mathbb{C}} = \mathcal{B}_{i,4}^{+} \cup \mathcal{B}_{i,4} \cup \mathcal{B}_{i,4}^{-}$, $\mathcal{O}_{i,3}^{\mathbb{C}} = \mathcal{B}_{i,3r}^{+} \cup \mathcal{B}_{i,3} \cup \mathcal{B}_{i,3}^{-} \cup \mathcal{B}_{i,3l}^{+}$, $\mathcal{B}_{i,3} = \mathcal{B}_{i,3r} \cup \{\mathbf{C}\} \cup \mathcal{B}_{i,3l}$ and $\mathcal{B}_{i,3r} \cap \mathcal{B}_{i,3l} = \varnothing$ for $i = 1, 2$.

Let $\mathcal{B} := \cup_{i=1}^{2} \cup_{j=3}^{4} \mathcal{B}_{i,j}$. Then $\theta(\hat{\boldsymbol{\omega}}, \mathbf{A}) = 0$ if and only if $\mathbf{A} \in \mathcal{B}$ for the fixed equation $\hat{\boldsymbol{\omega}}$, where θ is defined by (2.5).

Note that there are two real parameters in each $\mathcal{O}_{i,j}^{\mathbb{C}}$, $1 \leq i \leq 2$, $3 \leq j \leq 4$. Now we recall the monotonicity of the continuous eigenvalue branches with respect to the two real parameters, which was obtained in [22].

Lemma 4.1 [22, Theorem 4.6]. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then, in each of the coordinate systems $\mathcal{O}_{1,3}^{\mathbb{C}}$, $\mathcal{O}_{1,4}^{\mathbb{C}}$, $\mathcal{O}_{2,3}^{\mathbb{C}}$ and $\mathcal{O}_{2,4}^{\mathbb{C}}$ in $\mathcal{B}^{\mathbb{C}}$, every continuous eigenvalue branch is always non-decreasing in the two real axis directions.*

For example, in $\mathcal{O}_{1,4}^{\mathbb{C}}$, every continuous eigenvalue branch is always non-decreasing in the a_{12}-direction and in the b_{21}-direction.

Theorem 4.1. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then for each $\mathbf{A} \in \mathcal{B}_{1,4}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues and for each $\mathbf{A} \in \mathcal{O}_{1,4}^{\mathbb{C}} \backslash \mathcal{B}_{1,4}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues $\lambda_n(\mathbf{A}) := \lambda_n(\hat{\boldsymbol{\omega}}, \mathbf{A})$, $0 \leq n \leq N-1$, which satisfy that*

(i) *$\lambda_n(\mathbf{A})$ are continuous in $\mathcal{O}_{1,4}^{\mathbb{C}} \backslash \mathcal{B}_{1,4}$;*

(ii) *$\lambda_n(\mathbf{A})$ restricted in $\mathcal{B}_{1,4}^{+}$ and $\mathcal{B}_{1,4}^{-}$ are always non-decreasing in the a_{12}-direction and in the b_{21}-direction;*

(iii) *$\lambda_n(\mathbf{A})$ are not continuous at each point of $\mathcal{B}_{1,4}$ and have the following asymptotic behaviors near any given $\mathbf{A}_0 \in \mathcal{B}_{1,4}$:*

$$\lim_{\mathcal{B}_{1,4}^{-} \cup \mathcal{B}_{1,4} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{A}_0),\ 0 \leq n \leq N-2,\ \lim_{\mathcal{B}_{1,4}^{-} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_{N-1}(\mathbf{A}) = +\infty, \quad (4.1)$$

$$\lim_{\mathcal{B}_{1,4}^{+} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_0(\mathbf{A}) = -\infty, \quad \lim_{\mathcal{B}_{1,4}^{+} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{A}_0),\ \ 1 \leq n \leq N-1. \quad (4.2)$$

Consequently, $\lambda_n(\mathbf{A})$ restricted in $\mathcal{B}_{1,4}^{-} \cup \mathcal{B}_{1,4}$ is continuous for each $0 \leq n \leq N-2$.

Proof. By (2.4), $r(\hat{\boldsymbol{\omega}}, \mathbf{A}) = 1$ for each $\mathbf{A} \in \mathcal{B}_{1,4}$, and $r(\hat{\boldsymbol{\omega}}, \mathbf{A}) = 2$ for each $\mathbf{A} \in \mathcal{O}^{\mathbb{C}}_{1,4} \backslash \mathcal{B}_{1,4}$. It follows from Lemma 2.4 that $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues for each $\mathbf{A} \in \mathcal{B}_{1,4}$ and $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues for each $\mathbf{A} \in \mathcal{O}^{\mathbb{C}}_{1,4} \backslash \mathcal{B}_{1,4}$.

Now, we show that (i) and (ii) hold. Since $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues $\lambda_n(\mathbf{A})$, $0 \le n \le N-1$, for each $\mathbf{A} \in \mathcal{B}^{+}_{1,4}$ (or $\mathcal{B}^{-}_{1,4}$), which is an open and connected subset of $\mathcal{B}^{\mathbb{C}}$, $\lambda_n(\mathbf{A})$ is continuous and locally a continuous eigenvalue branch in $\mathcal{B}^{+}_{1,4}$ (or $\mathcal{B}^{-}_{1,4}$) for each $0 \le n \le N-1$ by Theorem 2.1. This, together with Lemma 4.1, implies its monotonicity in $\mathcal{B}^{+}_{1,4}$ (or $\mathcal{B}^{-}_{1,4}$).

Next, we show that (iii) holds. It suffices to show that (4.1)–(4.2) hold for any given $\mathbf{A}_0 \in \mathcal{B}_{1,4}$. Let (r_1, r_2) be a finite interval such that $\lambda_j(\mathbf{A}_0) \in (r_1, r_2)$ for all $0 \le j \le N-2$. By Lemma 2.5, there exists a neighborhood $\mathcal{V}$ of $\mathbf{A}_0$ in $\mathcal{O}^{\mathbb{C}}_{1,4}$ such that for each $\mathbf{A} \in \mathcal{V}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues in $[r_1, r_2]$ that are all in (r_1, r_2). Let

$$\mathcal{V}^{-} := \mathcal{B}^{-}_{1,4} \cap \mathcal{V},\ \mathcal{V}^{0} := \mathcal{B}_{1,4} \cap \mathcal{V},\ \mathcal{V}^{+} := \mathcal{B}^{+}_{1,4} \cap \mathcal{V}.$$

Note that $\mathcal{V}$ can be chosen such that $\mathcal{V}^{-}$, $\mathcal{V}^{0}$ and $\mathcal{V}^{+}$ are connected. Since $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues for each $\mathbf{A} \in \mathcal{V}^{-}$, it has exactly an eigenvalue, denoted by $\hat{\lambda}(\mathbf{A})$, outside $[r_1, r_2]$. By (i) of Theorem 2.3, either $G := \{\hat{\lambda}(\mathbf{A}) : \mathbf{A} \in \mathcal{V}^{-}\} \subset (-\infty, r_1)$ or $G \subset (r_2, +\infty)$.

Suppose that

$$\mathbf{A}_0 = \begin{bmatrix} 1 & 1/\hat{f}_0 & \bar{z}^0 & 0 \\ 0 & z^0 & b^0_{21} & 1 \end{bmatrix} \in \mathcal{B}_{1,4}.$$

Denote

$$\mathbf{A}(s) := \begin{bmatrix} 1 & s & \bar{z}^0 & 0 \\ 0 & z^0 & b^0_{21} & 1 \end{bmatrix},\ s \in \mathbb{R}.$$

Then $\mathbf{A}_0 = \mathbf{A}(1/\hat{f}_0)$. Note that each $(\hat{\boldsymbol{\omega}}, \mathbf{A}(s))$ has exactly N eigenvalues for each $s \in (-\infty, 1/\hat{f}_0)$, and $(\hat{\boldsymbol{\omega}}, \mathbf{A}(1/\hat{f}_0))$ has exactly $N-1$ eigenvalues. By (ii), $\lambda_n(\mathbf{A}(s))$ is non-decreasing in $(-\infty, 1/\hat{f}_0)$ for each $0 \le n \le N-1$. Hence, $\lim_{s \to 1/\hat{f}_0^-} \lambda_{N-1}(\mathbf{A}(s)) = +\infty$ by (ii) of Lemma 2.7. This implies that there exists an $\mathbf{A}_1 \in \mathcal{V}^{-}$ such that $\lambda_{N-1}(\mathbf{A}_1) > r_2$. Hence, $\hat{\lambda}(\mathbf{A}_1) = \lambda_{N-1}(\mathbf{A}_1)$. Thus, again by (i) of Theorem 2.3, $G = \{\lambda_{N-1}(\mathbf{A}) : \mathbf{A} \in \mathcal{V}^{-}\} \subset (r_2, +\infty)$. By (ii) of Theorem 2.3, the second relation in (4.1) holds.

With a similar argument to the proof of the second relation in (4.1), one can show that the first relation in (4.2) holds.

Since $\mathcal{V}^0$ is connected and $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues for each $\mathbf{A} \in \mathcal{V}^0$, by Theorem 2.1 λ_n restricted in $\mathcal{V}^0$ is continuous and locally a continuous eigenvalue branch for each $0 \le n \le N-2$. This, together with Theorem 2.2, shows that the first relation in (4.1) and the second relation in (4.2) hold. This completes the proof.

Theorem 4.2. *Fix a difference equation* $\hat{\boldsymbol{\omega}}$. *Then similar results in Theorem* 4.1 *hold, where* $\mathcal{O}^{\mathbb{C}}_{1,4}$, $\mathcal{B}_{1,4}$, $\mathcal{B}^{+}_{1,4}$, $\mathcal{B}^{-}_{1,4}$, *and* a_{12} *are replaced by* $\mathcal{O}^{\mathbb{C}}_{2,4}$, $\mathcal{B}_{2,4}$, $\mathcal{B}^{+}_{2,4}$, $\mathcal{B}^{-}_{2,4}$, *and* a_{11}, *respectively. The corresponding relations in* (4.1)–(4.2) *are denoted by* (4.1′)–(4.2′).

Proof. Since the proof is similar to that of Theorem 4.1, we omit its details.

Theorem 4.3. *Fix a difference equation* $\hat{\boldsymbol{\omega}}$*. Then* $(\hat{\boldsymbol{\omega}}, \mathbf{C})$ *has exactly* $N-2$ *eigenvalues, and for each* $\mathbf{A} \in \mathcal{B}_{1,3} \backslash \{\mathbf{C}\}$*,* $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ *has exactly* $N-1$ *eigenvalues, and for each* $\mathbf{A} \in \mathcal{O}_{1,3}^{\mathbb{C}} \backslash \mathcal{B}_{1,3}$*,* $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ *has exactly* N *eigenvalues* $\lambda_n(\mathbf{A})$*,* $0 \leq n \leq N-1$*, which satisfy that*

(i) $\lambda_n(\mathbf{A})$ *are continuous in* $\mathcal{O}_{1,3}^{\mathbb{C}} \backslash \mathcal{B}_{1,3}$*;*

(ii) $\lambda_n(\mathbf{A})$ *restricted in* $\mathcal{B}_{1,3}^-$*,* $\mathcal{B}_{1,3r}^+$ *and* $\mathcal{B}_{1,3l}^+$ *are non-decreasing in the* a_{12}*-direction and in the* b_{22}*-direction;*

(iii) $\lambda_n(\mathbf{A})$ *are not continuous at each point of* $\mathcal{B}_{1,3}$ *and furthermore,*

(iiia) *they have the following asymptotic behaviors near any given* $\mathbf{A}_0 \in \mathcal{B}_{1,3r}$*:*

$$\begin{gathered}
\lim_{\mathcal{B}_{1,3}^- \cup \mathcal{B}_{1,3r} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{A}_0), 0 \leq n \leq N-2, \\
\lim_{\mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{A}_0} \lambda_{N-1}(\mathbf{A}) = +\infty, \quad \lim_{\mathcal{B}_{1,3r}^+ \ni \mathbf{A} \to \mathbf{A}_0} \lambda_0(\mathbf{A}) = -\infty, \\
\lim_{\mathcal{B}_{1,3r}^+ \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{A}_0), \ 1 \leq n \leq N-1;
\end{gathered} \tag{4.3}$$

(iiib) *they have the following asymptotic behaviors near any given* $\mathbf{A}_0 \in \mathcal{B}_{1,3l}$*:*

$$\begin{gathered}
\lim_{\mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{A}_0} \lambda_0(\mathbf{A}) = -\infty, \\
\lim_{\mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{A}_0), \ 1 \leq n \leq N-1, \\
\lim_{\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3l} \ni \mathbf{A} \to \mathbf{A}_0} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{A}_0), \ \ 0 \leq n \leq N-2, \\
\lim_{\mathcal{B}_{1,3l}^+ \ni \mathbf{A} \to \mathbf{A}_0} \lambda_{N-1}(\mathbf{A}) = +\infty;
\end{gathered} \tag{4.4}$$

(iiic) *they have the following asymptotic behaviors near* $\mathbf{C}$*:*

$$\begin{gathered}
\lim_{\mathcal{B}_{1,3r}^+ \cup \mathcal{B}_{1,3r} \cup \mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{C}} \lambda_0(\mathbf{A}) = -\infty, \quad \lim_{\mathcal{B}_{1,3r}^+ \ni \mathbf{A} \to \mathbf{C}} \lambda_1(\mathbf{A}) = -\infty, \\
\lim_{\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{C}), \ 0 \leq n \leq N-3, \\
\lim_{\mathcal{B}_{1,3r} \cup \mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{C}), \ 1 \leq n \leq N-2, \\
\lim_{\mathcal{B}_{1,3r}^+ \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_{n-2}(\mathbf{C}), \ 2 \leq n \leq N-1, \\
\lim_{\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_{N-2}(\mathbf{A}) = +\infty, \quad \lim_{\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3}^- \ni \mathbf{A} \to \mathbf{C}} \lambda_{N-1}(\mathbf{A}) = +\infty.
\end{gathered} \tag{4.5}$$

And consequently, $\lambda_n(\mathbf{A})$ *restricted in* $\mathcal{B}_{1,3}^- \cup \mathcal{B}_{1,3r}$ *and* $\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3l} \cup \{\mathbf{C}\}$ *is continuous for each* $0 \leq n \leq N-3$*, and* $\lambda_{N-2}(\mathbf{A})$ *restricted in* $\mathcal{B}_{1,3}^- \cup \mathcal{B}_{1,3r}$ *and* $\mathcal{B}_{1,3l}^+ \cup \mathcal{B}_{1,3l}$ *is continuous.*

Proof. Since the proofs of (i), (ii), (iiia) and (iiib) are similar to those of Theorem 4.1, we omit their details.

The rest is to show that (iiic) holds. Note that $(\hat{\boldsymbol{\omega}}, \mathbf{C})$ has exactly $N-2$ eigenvalues. Let (r_1, r_2) be a finite interval such that $\lambda_j(\mathbf{C}) \in (r_1, r_2)$ for all $0 \leq j \leq N-3$. By Lemma

2.5, there exists a neighborhood $\mathcal{V}_1$ of $\mathbf{C}$ in $\mathcal{O}^{\mathbb{C}}_{1,3}$ such that for each $\mathbf{A} \in \mathcal{V}_1$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-2$ eigenvalues in $[r_1, r_2]$ that are all in (r_1, r_2). Note that $\mathcal{V}_1$ can be chosen such that $\mathcal{V}_1 \cap \mathcal{B}^+_{1,3r}$, $\mathcal{V}_1 \cap \mathcal{B}_{1,3r}$, $\mathcal{V}_1 \cap \mathcal{B}^-_{1,3}$, $\mathcal{V}_1 \cap \mathcal{B}_{1,3l}$, and $\mathcal{V}_1 \cap \mathcal{B}^+_{1,3l}$ are connected. Then we divide our proof in three steps.

Step 1. We show that

$$\lim_{\mathcal{B}_{1,3r} \ni \mathbf{A} \to \mathbf{C}} \lambda_0(\mathbf{A}) = -\infty, \lim_{\mathcal{B}_{1,3r} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{C}),\ 1 \le n \le N-2, \tag{4.6}$$

$$\lim_{\mathcal{B}_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{C}), 0 \le n \le N-3, \lim_{\mathcal{B}_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_{N-2}(\mathbf{A}) = +\infty. \tag{4.7}$$

Since for each $\mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}_{1,3r}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues, and then has exactly an eigenvalue, denoted by $\hat{\lambda}(\mathbf{A})$, outside $[r_1, r_2]$. By (i) of Theorem 2.3, $H := \{\hat{\lambda}(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}_{1,3r}\} \subset (-\infty, r_1)$ or $H \subset (r_2, +\infty)$.

Let

$$\mathbf{A}_1(s) = \begin{bmatrix} 1 & s & 0 & 0 \\ 0 & 0 & -1 & 0 \end{bmatrix},\ s \in \mathbb{R}.$$

Then $\mathbf{A}_1(1/\hat{f}_0) = \mathbf{C}$. For each $s \in (1/\hat{f}_0, +\infty)$, $(\hat{\boldsymbol{\omega}}, \mathbf{A}_1(s))$ has exactly $N-1$ eigenvalues and $(\hat{\boldsymbol{\omega}}, \mathbf{A}_1(1/\hat{f}_0))$ has exactly $N-2$ eigenvalues. By Theorem 2.1, $\lambda_n(\mathbf{A}_1(s))$ is continuous and locally a continuous eigenvalue branch in $(1/\hat{f}_0, +\infty)$ for each $0 \le n \le N-2$. This, together with Lemma 4.1, implies that $\lambda_n(\mathbf{A}_1(s))$ is non-decreasing in $(1/\hat{f}_0, +\infty)$ for each $0 \le n \le N-2$. Hence, $\lim_{s \to 1/\hat{f}_0^+} \lambda_0(\mathbf{A}_1(s)) = -\infty$ by (iv) of Lemma 2.7. This implies that there exists an $\mathbf{A}_1 \in \mathcal{V}_1 \cap \mathcal{B}_{1,3r}$ such that $\lambda_0(\mathbf{A}_1) < r_1$. Hence, $\hat{\lambda}(\mathbf{A}_1) = \lambda_0(\mathbf{A}_1)$. Thus, again by (i) of Theorem 2.3, $H = \{\lambda_0(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}_{1,3r}\} \subset (-\infty, r_1)$. Then it follows from (ii) of Theorem 2.3 that the first relation in (4.6) holds.

With a similar argument to the proof of the first relation in (4.6), one can show that the second relation in (4.7) holds.

It follows from Theorem 2.2 that the second relation in (4.6) and the first relation in (4.7) hold.

Step 2. We show that

$$\lim_{\mathcal{B}^+_{1,3r} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = -\infty,\ n = 0, 1, \quad \lim_{\mathcal{B}^+_{1,3r} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_{n-2}(\mathbf{C}),\ \ 2 \le n \le N-1, \tag{4.8}$$

$$\lim_{\mathcal{B}^+_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{C}), 0 \le n \le N-3, \quad \lim_{\mathcal{B}^+_{1,3l} \ni \mathbf{A} \to \mathbf{C}} \lambda_n(\mathbf{A}) = +\infty,\ n = N-2, N-1. \tag{4.9}$$

Since for each $\mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^+_{1,3r}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues, and then has exactly two eigenvalues, denoted by $\hat{\lambda}_1(\mathbf{A}) \le \hat{\lambda}_2(\mathbf{A})$, outside $[r_1, r_2]$. By (i) of Theorem 2.3, either $I_n := \{\hat{\lambda}_n(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^+_{1,3r}\} \subset (-\infty, r_1)$ or $I_n \subset (r_2, +\infty)$ for each $n = 1, 2$.

Let

$$\mathbf{A}_2(s) = \begin{bmatrix} 1 & s + 1/\hat{f}_0 & 0 & 0 \\ 0 & 0 & -1 & s \end{bmatrix},\ s \in \mathbb{R}.$$

Then $\mathbf{A}_2(0) = \mathbf{C}$. $(\hat{\boldsymbol{\omega}},\ \mathbf{A}_2(s))$ has exactly N eigenvalues for each $s \in (0, +\infty)$, and $(\hat{\boldsymbol{\omega}}, \mathbf{A}_2(0))$ has exactly $N-2$ eigenvalues. By (ii), $\lambda_n(\mathbf{A}_2(s))$ is non-decreasing in $(0, +\infty)$

for each $0 \le n \le N-1$. Hence, by (iv) of Lemma 2.7, $\lim_{s\to 0^+} \lambda_n(\mathbf{A}_2(s)) = -\infty, n = 0, 1$. This implies that there exists $\mathbf{A}_2 \in \mathcal{V}_1 \cap \mathcal{B}^+_{1,3r}$ such that $\lambda_n(\mathbf{A}_2) < r_1$, $n = 0, 1$. Hence, $\hat{\lambda}_n(\mathbf{A}_2) = \lambda_{n-1}(\mathbf{A}_2)$, $n = 1, 2$. Thus, again by (i) of Theorem 2.3, $I_n = \{\lambda_{n-1}(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^+_{1,3r}\} \subset (-\infty, r_1)$, $n = 1, 2$. By (ii) of Theorem 2.3, the first relation in (4.8) holds.

With a similar argument to the proof of the first relation in (4.8), one can show that the second relation in (4.9) holds.

It follows from Theorem 2.2 that the second relation in (4.8) and the first relation in (4.9) hold.

Step 3. We show that

$$\lim_{\mathcal{B}^-_{1,3}\ni\mathbf{A}\to\mathbf{C}} \lambda_0(\mathbf{A}) = -\infty, \quad \lim_{\mathcal{B}^-_{1,3}\ni\mathbf{A}\to\mathbf{C}} \lambda_{N-1}(\mathbf{A}) = +\infty, \tag{4.10}$$

$$\lim_{\mathcal{B}^-_{1,3}\ni\mathbf{A}\to\mathbf{C}} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{C}), \quad 1 \le N \le N-2. \tag{4.11}$$

Since for each $\mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^-_{1,3}$, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly N eigenvalues, and then has exactly two eigenvalues, denoted by $\lambda'_1(\mathbf{A}) \le \lambda'_2(\mathbf{A})$, outside $[r_1, r_2]$. By (i) of Theorem 2.3, either $J_n := \{\lambda'_n(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^-_{1,3}\} \subset (-\infty, r_1)$ or $J_n \subset (r_2, +\infty)$ for each $n = 1, 2$. Let $\mathbf{A}_3 \in \mathcal{V}_1 \cap \mathcal{B}_{1,3r}$ and $\mathbf{A}_4 \in \mathcal{V}_1 \cap \mathcal{B}_{1,3l}$. Then it follows from Step 1 that $\lambda_0(\mathbf{A}_3) \in (-\infty, r_1)$ and $\lambda_{N-2}(\mathbf{A}_4) \in (r_2, +\infty)$. By Lemma 2.6, $\lambda_0(\mathbf{A}_3)$ lies in a continuous eigenvalue branch Λ_1 defined in a connected neighborhood $\mathcal{V}_2$ of $\mathbf{A}_3$ in $\mathcal{O}^{\mathbb{C}}_{1,3}$. Thus, one can choose $\mathbf{A}_5 \in \mathcal{V}_1 \cap \mathcal{V}_2 \cap \mathcal{B}^-_{1,3}$ sufficiently close to $\mathbf{A}_3$ such that $\Lambda_1(\mathbf{A}_5) \in (-\infty, r_1)$. Similarly, $\lambda_{N-2}(\mathbf{A}_4)$ lies in a continuous eigenvalue branch Λ_2 defined in a connected neighborhood $\mathcal{V}_3$ of $\mathbf{A}_4$ in $\mathcal{O}^{\mathbb{C}}_{1,3}$. Thus, one can choose $\mathbf{A}_6 \in \mathcal{V}_1 \cap \mathcal{V}_3 \cap \mathcal{B}^-_{1,3}$ sufficiently close to $\mathbf{A}_4$ such that $\Lambda_2(\mathbf{A}_6) \in (r_2, +\infty)$. Since either $J_n \subset (-\infty, r_1)$ or $J_n \subset (r_2, +\infty)$ for each $n = 1, 2$, again by (i) of Theorem 2.3, $J_1 = \{\lambda_0(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^-_{1,3}\} \subset (-\infty, r_1)$ and $J_2 = \{\lambda_{N-1}(\mathbf{A}) : \mathbf{A} \in \mathcal{V}_1 \cap \mathcal{B}^-_{1,3}\} \subset (r_2, +\infty)$. By (ii) of Theorem 2.3, (4.10) holds. With a similar argument to that used in the proof of Theorem 2.2, one can show that (4.11) holds. The whole proof is complete.

Theorem 4.4. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then similar results in Theorem* 4.3 *hold, where $\mathcal{O}^{\mathbb{C}}_{1,3}$, $\mathcal{B}_{1,3}$, $\mathcal{B}_{1,3r}$, $\mathcal{B}_{1,3l}$, $\mathcal{B}^-_{1,3}$, $\mathcal{B}^+_{1,3r}$, $\mathcal{B}^+_{1,3l}$, and a_{12} are replaced by $\mathcal{O}^{\mathbb{C}}_{2,3}$, $\mathcal{B}_{2,3}$, $\mathcal{B}_{2,3r}$, $\mathcal{B}_{2,3l}$, $\mathcal{B}^-_{2,3}$, $\mathcal{B}^+_{2,3r}$, $\mathcal{B}^+_{2,3l}$, and a_{11}, respectively. The corresponding relations in* (4.3)–(4.5) *are denoted by* (4.3′)–(4.5′).

Proof. Since the proof is similar to that of Theorem 4.3, we omit its details.

Combining Theorems 4.1–4.4 yields the continuity and discontinuity sets for each of the n-th eigenvalue function in $\mathcal{B}^{\mathbb{C}}$:

Theorem 4.5. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then $(\hat{\boldsymbol{\omega}}, \mathbf{C})$ has exactly $N-2$ eigenvalues, $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly $N-1$ eigenvalues for each $\mathbf{A} \in \mathcal{B}\backslash\{\mathbf{C}\}$, and $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ has exactly*

N eigenvalues for each $\mathbf{A} \in \mathcal{B}^{\mathbb{C}} \backslash \mathcal{B}$. Moreover, the n-th eigenvalue function $\lambda_n(\mathbf{A})$ is continuous in $\mathcal{B}^{\mathbb{C}} \backslash \mathcal{B}$ and not continuous at each point of $\mathcal{B}$ for each $0 \leq n \leq N-1$.

Now we apply Theorems 4.1–4.4 to the separated and coupled BCs, respectively. We introduce the following notations for convenience: $\xi := \arctan(-1/\hat{f}_0) + \pi$ for $\hat{f}_0 > 0$, and $\xi := \arctan(-1/\hat{f}_0)$ for $\hat{f}_0 < 0$;

$$\begin{aligned}
\mathcal{B}_{S_1} &:= \{\mathbf{S}_{\alpha,\beta} \in \mathcal{B}_S : (\alpha,\beta) \in \{\xi\} \times (0,\pi] \text{ or } (\alpha,\beta) \in [0,\pi) \times \{\pi\}\}, \\
\mathcal{B}_{C_1} &:= \{[e^{i\gamma}K \mid -I] \in \mathcal{B}_C : \gamma \in [0,\pi),\ \ K \in SL(2,\mathbb{R}),\ k_{12} \neq 0,\ k_{11}/k_{12} = \hat{f}_0\}, \\
\mathcal{B}_{C_1}^{+} &:= \{[e^{i\gamma}K \mid -I] \in \mathcal{B}_C : \gamma \in [0,\pi),\ \ K \in SL(2,\mathbb{R}),\ k_{12} \neq 0,\ k_{11}/k_{12} > \hat{f}_0\}, \\
\mathcal{B}_{C_1}^{-} &:= \{[e^{i\gamma}K \mid -I] \in \mathcal{B}_C : \gamma \in [0,\pi),\ \ K \in SL(2,\mathbb{R}),\ k_{12} \neq 0,\ k_{11}/k_{12} < \hat{f}_0\},
\end{aligned}$$

where $\mathbf{S}_{\alpha,\beta}$ is defined by (2.2). By Lemma 2.2, $\theta(\hat{\boldsymbol{\omega}}, \mathbf{A}) = 0$ if and only if $\mathbf{A} \in \mathcal{B}_{S_1} \cup \mathcal{B}_{C_1}$, where θ is defined by (2.5). Thus

$$\mathcal{B} = \mathcal{B}_{S_1} \cup \mathcal{B}_{C_1}. \tag{4.12}$$

The following result gives the continuity and discontinuity sets of λ_n restricted in $\mathcal{B}_S$ and asymptotic behaviors of λ_n in $\mathcal{B}^{\mathbb{C}}$ near each discontinuity point.

Corollary 4.1. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then $(\hat{\boldsymbol{\omega}}, \mathbf{S}_{\xi,\pi})$ has exactly $N-2$ eigenvalues, and $(\hat{\boldsymbol{\omega}}, \mathbf{S}_{\alpha,\beta})$ has exactly $N-1$ eigenvalues for each $\mathbf{S}_{\alpha,\beta} \in \mathcal{B}_{S_1} \backslash \{\mathbf{S}_{\xi,\pi}\}$, and $(\hat{\boldsymbol{\omega}}, \mathbf{S}_{\alpha,\beta})$ has exactly N eigenvalues for each $\mathbf{S}_{\alpha,\beta} \in \mathcal{B}_S \backslash \mathcal{B}_{S_1}$. Moreover, the n-th eigenvalue function λ_n in $\mathcal{B}^{\mathbb{C}}$ is continuous in $\mathcal{B}_S \backslash \mathcal{B}_{S_1}$ and not continuous at each point of $\mathcal{B}_{S_1}$ for each $0 \leq n \leq N-1$, and furthermore,*

- (i) *for any fixed $\beta_0 \in (0,\pi)$, (4.1)–(4.2) and (4.1′)–(4.2′) hold for $\mathbf{A}_0$ replaced by $\mathbf{S}_{\xi,\beta_0}$;*
- (ii) *for any fixed $\alpha_0 \in [0,\pi) \backslash \{\xi, \pi/2\}$, (4.3) holds for $\mathbf{A}_0$ replaced by $\mathbf{S}_{\alpha_0,\pi}$ in the case that $\mathbf{S}_{\alpha_0,\pi} \in \mathcal{B}_{1,3r}$, and (4.4) holds for $\mathbf{A}_0$ replaced by $\mathbf{S}_{\alpha_0,\pi}$ in the other case that $\mathbf{S}_{\alpha_0,\pi} \in \mathcal{B}_{1,3l}$;*
- (iii) *(4.3′) holds for $\mathbf{A}_0$ replaced by $\mathbf{S}_{\pi/2,\pi}$ in the case that $\mathbf{S}_{\pi/2,\pi} \in \mathcal{B}_{2,3r}$, and (4.4′) holds for $\mathbf{A}_0$ replaced by $\mathbf{S}_{\pi/2,\pi}$ in the other case that $\mathbf{S}_{\pi/2,\pi} \in \mathcal{B}_{2,3l}$;*
- (iv) *(4.5) and (4.5′) hold for $\mathbf{C}$ replaced by $\mathbf{S}_{\xi,\pi}$.*

Proof. The number of eigenvalues of each $(\hat{\boldsymbol{\omega}}, \mathbf{S}_{\alpha,\beta})$ can be easily verified by Lemma 2.4. Since $\mathcal{B}_S \backslash \mathcal{B}_{S_1} \subset \mathcal{B}^{\mathbb{C}} \backslash \mathcal{B}$, λ_n in $\mathcal{B}^{\mathbb{C}}$ is continuous in $\mathcal{B}_S \backslash \mathcal{B}_{S_1}$ for each $0 \leq n \leq N-1$ by Theorem 4.5. Then, it suffices to show that (i)–(iv) hold.

(i) Fix any $\beta_0 \in (0,\pi)$. It is clear that

$$\mathbf{S}_{\xi,\beta_0} = \begin{bmatrix} 1 & 1/\hat{f}_0 & 0 & 0 \\ 0 & 0 & -\cot\beta_0 & 1 \end{bmatrix} \in \mathcal{B}_{1,4} \cap \mathcal{B}_{2,4}.$$

Applying Theorems 4.1-4.2 to $\mathbf{S}_{\xi,\beta_0}$, one gets that (i) holds.

(ii) Fix any $\alpha_0 \in [0,\pi)\backslash\{\xi,\pi/2\}$. It is clear that

$$\mathbf{S}_{\alpha_0,\pi} = \left[\begin{array}{cccc} 1 & -\tan\alpha_0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array}\right] \in \mathcal{B}_{1,3r} \cup \mathcal{B}_{1,3l}.$$

Applying (iiia)–(iiib) in Theorem 4.3 to $\mathbf{S}_{\alpha_0,\pi}$, one gets that (ii) holds.

(iii) It is clear that

$$\mathbf{S}_{\pi/2,\pi} = \left[\begin{array}{cccc} 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array}\right] \in \mathcal{B}_{2,3r} \cup \mathcal{B}_{2,3l}.$$

Applying (iiia)–(iiib) in Theorem 4.4 to $\mathbf{S}_{\pi/2,\pi}$, one gets that (iii) holds.

(iv) It is clear that

$$\mathbf{S}_{\xi,\pi} = \left[\begin{array}{cccc} 1 & 1/\hat{f}_0 & 0 & 0 \\ 0 & 0 & -1 & 0 \end{array}\right] = \mathbf{C}.$$

Applying (iiic) in Theorems 4.3–4.4 to $\mathbf{S}_{\xi,\pi}$, one gets that (iv) holds. The proof is complete.

The following lemma is the monotonicity result of continuous eigenvalue branches with respect to the two real parameters α and β for $\mathbf{S}_{\alpha,\beta}$.

Lemma 4.2 [22, Theorem 4.4]. *Each continuous eigenvalue branch over $\mathcal{B}_S$ is always strictly decreasing in the α-direction and always strictly increasing in the β-direction.*

Now, we consider λ_n restricted in $\mathcal{B}_S$ for each $0 \le n \le N-1$. For a fixed $\beta_0 \in (0,\pi]$, set $\lambda_n(\alpha) := \lambda_n(\mathbf{S}_{\alpha,\beta_0})$, and for a fixed $\alpha_0 \in [0,\pi)$, set $\lambda_n(\beta) := \lambda_n(\mathbf{S}_{\alpha_0,\beta})$ for convenience.

Corollary 4.2. *Fix a difference equation $\hat{\boldsymbol{\omega}}$. Then*

(i) *for any fixed $\beta_0 \in (0,\pi)$, the n-th eigenvalue functions $\lambda_n(\alpha)$ are strictly decreasing in $[0,\xi)$ or (ξ,π) for all $0 \le n \le N-1$, and have the following asymptotic behaviors near 0 and ξ:*

$$\lim_{\alpha\to\pi^-} \lambda_n(\alpha) = \lambda_n(0), \quad 0 \le n \le N-1, \tag{4.13}$$

$$\lim_{\alpha\to\xi^-} \lambda_0(\alpha) = -\infty, \quad \lim_{\alpha\to\xi^-} \lambda_n(\alpha) = \lambda_{n-1}(\xi), \quad 1 \le n \le N-1,$$

$$\lim_{\alpha\to\xi^+} \lambda_n(\alpha) = \lambda_n(\xi), \ 0 \le n \le N-2, \quad \lim_{\alpha\to\xi^+} \lambda_{N-1}(\alpha) = +\infty,$$

and consequently, $\lambda_n(\alpha)$ is continuous in $[\xi,\pi)$ for each $0 \le n \le N-2$;

(ii) *for any fixed $\alpha_0 \in [0,\pi)\backslash\{\xi\}$, the n-th eigenvalue functions $\lambda_n(\beta)$ are strictly increasing in $(0,\pi)$ for all $0 \le n \le N-1$, and have the following asymptotic behaviors near π:*

$$\lim_{\beta\to\pi^-} \lambda_n(\beta) = \lambda_n(\pi), \quad 0 \le n \le N-2, \quad \lim_{\beta\to\pi^-} \lambda_{N-1}(\beta) = +\infty,$$

$$\lim_{\beta\to 0^+} \lambda_0(\beta) = -\infty, \quad \lim_{\beta\to 0^+} \lambda_n(\beta) = \lambda_{n-1}(\pi), \quad 1 \le n \le N-1,$$

and consequently, $\lambda_n(\beta)$ is continuous in $(0,\pi]$ for each $0 \le n \le N-2$;

(iii) *for $\beta_0 = \pi$, similar results in* (i) *hold for $N-2$, $N-1$ replaced by $N-3$, $N-2$, respectively.*

(iv) *for* $\alpha_0 = \xi$, *similar results in* (ii) *hold for* $N-2$, $N-1$ *replaced by* $N-3$, $N-2$, *respectively.*

Proof. It suffices to show that (4.13) holds since the rest is direct consequence of Theorem 2.1, Lemma 2.7, Corollary 4.1 and Lemma 4.2.

Fixed $\beta_0 \in (0, \pi)$. Since $\{\mathbf{S}_{\alpha,\beta_0} : \alpha \in [0, \xi) \cup (\xi, \pi)\}$ is connected and $(\hat{\boldsymbol{\omega}}, \mathbf{S}_{\alpha,\beta_0})$ has exactly N eigenvalues for each $\alpha \in [0, \xi) \cup (\xi, \pi)$, the n-th eigenvalue function λ_n is continuous and locally a continuous eigenvalue branch in $\{\mathbf{S}_{\alpha,\beta_0} : \alpha \in [0, \xi) \cup (\xi, \pi)\}$ for each $0 \le n \le N-1$ by Theorem 2.1. Then $\lim_{\alpha \to \pi^-} \mathbf{S}_{\alpha,\beta_0} = \mathbf{S}_{0,\beta_0}$ implies that (4.13) holds. This completes the proof.

The following result gives the continuity and discontinuity sets of λ_n restricted in $\mathcal{B}_C$ and asymptotic behaviors of λ_n in $\mathcal{B}^{\mathbb{C}}$ near each discontinuity point.

Corollary 4.3. *Fix a difference equation* $\hat{\boldsymbol{\omega}}$. *Then* $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ *has exactly* $N-1$ *eigenvalues for each* $\mathbf{A} \in \mathcal{B}_{C_1}$, *and* $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ *has exactly* N *eigenvalues for each* $\mathbf{A} \in \mathcal{B}_C \backslash \mathcal{B}_{C_1}$. *Moreover, the n-th eigenvalue function* λ_n *in* $\mathcal{B}^{\mathbb{C}}$ *is continuous in* $\mathcal{B}_C \backslash \mathcal{B}_{C_1}$, *and not continuous at each point of* $\mathcal{B}_{C_1}$ *for each* $0 \le n \le N-1$; *and for any given* $\mathbf{A}_1 \in \mathcal{B}_{C_1}$, (4.1)–(4.2) *and* (4.1′)–(4.2′) *hold for* $\mathbf{A}_0$ *replaced by* $\mathbf{A}_1$.

Proof. The number of eigenvalues of $(\hat{\boldsymbol{\omega}}, \mathbf{A})$ can be easily verified by Lemma 2.4 for each $\mathbf{A} \in \mathcal{B}_C$. Since $\mathcal{B}_C \backslash \mathcal{B}_{C_1} \subset \mathcal{B}^{\mathbb{C}} \backslash \mathcal{B}$, λ_n in $\mathcal{B}^{\mathbb{C}}$ is continuous in $\mathcal{B}_C \backslash \mathcal{B}_{C_1}$ for each $0 \le n \le N-1$ by Theorem 4.5. Let $\mathbf{A}_1 := [e^{i\gamma}K \,|\, -I] \in \mathcal{B}_{C_1}$. Then $k_{11} \neq 0$ and $k_{12} \neq 0$. By Lemma 3.18 in [13],

$$\mathbf{A}_1 = \begin{bmatrix} 1 & k_{12}/k_{11} & -e^{-i\gamma}/k_{11} & 0 \\ 0 & -e^{i\gamma}/k_{11} & -k_{21}/k_{11} & 1 \end{bmatrix} = \begin{bmatrix} -k_{11}/k_{12} & -1 & e^{-i\gamma}/k_{12} & 0 \\ e^{i\gamma}/k_{12} & 0 & -k_{22}/k_{12} & 1 \end{bmatrix} \in \mathcal{O}^{\mathbb{C}}_{1,4} \cap \mathcal{O}^{\mathbb{C}}_{2,4}.$$

Since $k_{11} - \hat{f}_0 k_{12} = 0$, $\mathbf{A}_1 \in \mathcal{B}_{1,4} \cap \mathcal{B}_{2,4}$. Hence, the conclusion holds by applying Theorems 4.1–4.2 to $\mathbf{A}_1$.

Let λ_n be restricted in $\mathcal{B}_C$ for each $0 \le n \le N-1$. The following result is a direct consequence of Corollary 4.3.

Corollary 4.4. *Fix a difference equation* $\hat{\boldsymbol{\omega}}$. *Then for any given* $\mathbf{A}_1 \in \mathcal{B}_{C_1}$, *the n-th eigenvalue functions* λ_n, $0 \le n \le N-1$, *have the following asymptotic behaviors near* $\mathbf{A}_1$:

$$\lim_{\mathcal{B}^-_{C_1} \ni \mathbf{A} \to \mathbf{A}_1} \lambda_0(\mathbf{A}) = -\infty, \quad \lim_{\mathcal{B}^-_{C_1} \ni \mathbf{A} \to \mathbf{A}_1} \lambda_n(\mathbf{A}) = \lambda_{n-1}(\mathbf{A}_1), \ \ 1 \le n \le N-1,$$
$$\lim_{\mathcal{B}^+_{C_1} \cup \mathcal{B}_{C_1} \ni \mathbf{A} \to \mathbf{A}_1} \lambda_n(\mathbf{A}) = \lambda_n(\mathbf{A}_1), \ \ 0 \le n \le N-2, \quad \lim_{\mathcal{B}^+_{C_1} \ni \mathbf{A} \to \mathbf{A}_1} \lambda_{N-1}(\mathbf{A}) = +\infty.$$

5. Continuity and discontinuity of the n-th eigenvalue function in the space of the SLPs

In this section, the continuous and discontinuous dependence of the n-th eigenvalue function on the SLP (1.1)–(1.2) is discussed. Its continuity and discontinuity sets in $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$ are given and its asymptotic behaviors near a discontinuity point are completely characterized.

Now, we introduce the following notations:

$$\mathcal{P}_{1,4} := \{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}} : a_{12} = 1/f_0\}, \mathcal{P}_{2,4} := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{2,4}^{\mathbb{C}} : a_{11} = -f_0\right\},$$

$$\mathcal{P}_{1,3} := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,3}^{\mathbb{C}} : (a_{12} - 1/f_0)b_{22} = |z|^2\right\},$$

$$\mathcal{P}_{2,3} := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{2,3}^{\mathbb{C}} : (a_{11} + f_0)b_{22} = |z|^2\right\},$$

$$\mathcal{P}_5 := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}} : \mathbf{A} = \begin{bmatrix} 1 & 1/f_0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}\right\},$$

$$\mathcal{P}_{1,3r} := \{(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{1,3} : a_{12} - 1/f_0 \geq 0, b_{22} \geq 0\} \backslash \mathcal{P}_5,$$

$$\mathcal{P}_{1,3l} := \{(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{1,3} : a_{12} - 1/f_0 \leq 0, b_{22} \leq 0\} \backslash \mathcal{P}_5,$$

$$\mathcal{P}_{1,3r}^+ := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,3}^{\mathbb{C}} : a_{12} - 1/f_0 \geq 0, b_{22} \geq 0, (a_{12} - 1/f_0)b_{22} > |z|^2\right\},$$

$$\mathcal{P}_{1,3l}^+ := \left\{(\boldsymbol{\omega}, \mathbf{A}) \in \Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,3}^{\mathbb{C}} : a_{12} - 1/f_0 \leq 0, b_{22} \leq 0, (a_{12} - 1/f_0)b_{22} > |z|^2\right\},$$

$$\mathcal{P}_{2,3r} := \{(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{2,3} : a_{11} + f_0 \geq 0, b_{22} \geq 0\} \backslash \mathcal{P}_5,$$

$$\mathcal{P}_{2,3l} := \{(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{2,3} : a_{11} + f_0 \leq 0, b_{22} \leq 0\} \backslash \mathcal{P}_5,$$

$\mathcal{P}_{1,4}^+$ and $\mathcal{P}_{2,4}^+$, and $\mathcal{P}_{1,4}^-$, $\mathcal{P}_{2,4}^-$, $\mathcal{P}_{1,3}^-$, and $\mathcal{P}_{2,3}^-$ are defined similarly as $\mathcal{E}^+$ and $\mathcal{E}^-$, respectively; $\mathcal{P}_{2,3r}^+$ and $\mathcal{P}_{2,3l}^+$ are defined similarly as $\mathcal{P}_{1,3r}^+$ and $\mathcal{P}_{1,3l}^+$, respectively. Then $\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{i,4}^{\mathbb{C}} = \mathcal{P}_{i,4}^+ \cup \mathcal{P}_{i,4} \cup \mathcal{P}_{i,4}^-$, $\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{i,3}^{\mathbb{C}} = \mathcal{P}_{i,3r}^+ \cup \mathcal{P}_{i,3} \cup \mathcal{P}_{i,3}^- \cup \mathcal{P}_{i,3l}^+$, $\mathcal{P}_{i,3} = \mathcal{P}_{i,3r} \cup \mathcal{P}_5 \cup \mathcal{P}_{i,3l}$ and $\mathcal{P}_{i,3r} \cap \mathcal{P}_{i,3l} = \varnothing$, where $i = 1, 2$.

Let $\mathcal{P} := \cup_{i=1}^2 \cup_{j=3}^4 \mathcal{P}_{i,j}$. Note that $\theta(\boldsymbol{\omega}, \mathbf{A}) = 0$ if and only if $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}$, where θ is defined by (2.5). Thus, $\mathcal{P}$ consists of the SLPs that have less than N eigenvalues.

Theorem 5.1. *Each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{1,4}$ has exactly $N-1$ eigenvalues and each $(\boldsymbol{\omega}, \mathbf{A}) \in \left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,4}$ has exactly N eigenvalues $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$, $0 \leq n \leq N-1$, which satisfy that*

(i) *$\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ are continuous in $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,4}$;*

(ii) *$\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ are not continuous at each point of $\mathcal{P}_{1,4}$ and have the following asymptotic behaviors near any given $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{P}_{1,4}$:*

$$\lim_{\mathcal{P}_{1,4}^- \cup \mathcal{P}_{1,4} \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_n(\boldsymbol{\omega}_0, \mathbf{A}_0), \quad 0 \leq n \leq N-2, \tag{5.1}$$

$$\lim_{\mathcal{P}_{1,4}^- \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_{N-1}(\boldsymbol{\omega}, \mathbf{A}) = +\infty, \quad \lim_{\mathcal{P}_{1,4}^+ \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_0(\boldsymbol{\omega}, \mathbf{A}) = -\infty, \tag{5.2}$$

$$\lim_{\mathcal{P}_{1,4}^+ \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n-1}(\boldsymbol{\omega}_0, \mathbf{A}_0), \quad 1 \leq n \leq N-1, \tag{5.3}$$

and consequently, λ_n restricted in $\mathcal{P}_{1,4}^- \cup \mathcal{P}_{1,4}$ is continuous for each $0 \leq n \leq N-2$.

Proof. By Lemma 2.4, each $(\boldsymbol{\omega}, \mathbf{A}) \in \left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,4}$ has exactly N eigenvalues, and each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{1,4}$ has exactly $N-1$ eigenvalues. Since $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,4}$ is an open subset of $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$, by Theorem 2.1 $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ is continuous in $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,4}$ for each $0 \le n \le N-1$.

The rest is to show that (5.1)–(5.3) hold for any given $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{P}_{1,4}$. Let (r_1, r_2) be a finite interval such that $\lambda_j(\boldsymbol{\omega}_0, \mathbf{A}_0) \in (r_1, r_2)$ for all $0 \le n \le N-2$. Then by Lemma 2.5 there exists a neighborhood $\mathcal{Y}$ of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in $\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,4}^{\mathbb{C}}$ such that each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{Y}$ has exactly $N-1$ eigenvalues in $[r_1, r_2]$ that are all in (r_1, r_2). Let

$$\mathcal{Y}^- := \mathcal{P}_{1,4}^- \cap \mathcal{Y},\ \mathcal{Y}^0 := \mathcal{P}_{1,4} \cap \mathcal{Y},\ \mathcal{Y}^+ := \mathcal{P}_{1,4}^+ \cap \mathcal{Y}.$$

Note that $\mathcal{Y}$ can be chosen such that $\mathcal{Y}^-$, $\mathcal{Y}^0$ and $\mathcal{Y}^+$ are connected. Since each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{Y}^+$ has exactly N eigenvalues, it has exactly one eigenvalue, denoted by $\hat{\lambda}(\boldsymbol{\omega}, \mathbf{A})$, outside $[r_1, r_2]$. By (i) of Theorem 2.3, either $L := \{\hat{\lambda}(\boldsymbol{\omega}, \mathbf{A}) : (\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{Y}^+\} \subset (-\infty, r_1)$ or $L \subset (r_2, +\infty)$. For the fix $\boldsymbol{\omega}_0$, there exists an $(\boldsymbol{\omega}_0, \mathbf{A}_1) \in \mathcal{Y}^+$ such that $\lambda_0(\boldsymbol{\omega}_0, \mathbf{A}_1) < r_1$ by the first relation in (4.2) in Theorem 4.1. Hence, $\hat{\lambda}(\boldsymbol{\omega}_0, \mathbf{A}_1) = \lambda_0(\boldsymbol{\omega}_0, \mathbf{A}_1)$. Thus, again by (i) of Theorem 2.3, $L = \{\lambda_0(\boldsymbol{\omega}, \mathbf{A}) : (\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{Y}^+\} \subset (-\infty, r_1)$. Then, by (ii) of Theorem 2.3, the second relation in (5.2) holds. With a similar argument, one can show that the first relation in (5.2) holds. Note that $\mathcal{Y}^0$ is connected and each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{Y}^0$ has exactly $N-1$ eigenvalues. So λ_n restricted in $\mathcal{Y}^0$ is continuous and locally a continuous eigenvalue branch for each $0 \le n \le N-2$ by Theorem 2.1. This, together with Theorem 2.2, yields that (5.1) and (5.3) hold. This completes the proof.

With a similar method used in the proof of Theorem 5.1, one can show that Theorems 5.2–5.4 hold.

Theorem 5.2. *Similar results in Theorem* 5.1 *hold for $\mathcal{O}_{1,4}^{\mathbb{C}}$, $\mathcal{P}_{1,4}$, $\mathcal{P}_{1,4}^+$ and $\mathcal{P}_{1,4}^-$ replaced by $\mathcal{O}_{2,4}^{\mathbb{C}}$, $\mathcal{P}_{2,4}$, $\mathcal{P}_{2,4}^+$ and $\mathcal{P}_{2,4}^-$, separately.*

Theorem 5.3. *Each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_5$ has exactly $N-2$ eigenvalues, and each $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_{1,3} \backslash \mathcal{P}_5$ has exactly $N-1$ eigenvalues, and each $(\boldsymbol{\omega}, \mathbf{A}) \in \left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,3}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,3}$ has exactly N eigenvalues $\lambda_n(\boldsymbol{\omega}, \mathbf{A})$, $0 \le n \le N-1$, which satisfy that*

- (i) *$\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ are continuous in $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{O}_{1,3}^{\mathbb{C}}\right) \backslash \mathcal{P}_{1,3}$;*
- (ii) *$\lambda_n(\boldsymbol{\omega}, \mathbf{A})$ are not continuous at each point of $\mathcal{P}_{1,3}$ and have the following asymptotic behaviors near any given $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{P}_{1,3r}$:*

$$\lim_{\mathcal{P}_{1,3}^- \cup \mathcal{P}_{1,3r} \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_n(\boldsymbol{\omega}_0, \mathbf{A}_0), \ \ 0 \le n \le N-2,$$
$$\lim_{\mathcal{P}_{1,3}^- \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_{N-1}(\boldsymbol{\omega}, \mathbf{A}) = +\infty, \quad \lim_{\mathcal{P}_{1,3r}^+ \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_0(\boldsymbol{\omega}, \mathbf{A}) = -\infty,$$
$$\lim_{\mathcal{P}_{1,3r}^+ \ni (\boldsymbol{\omega}, \mathbf{A}) \to (\boldsymbol{\omega}_0, \mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n-1}(\boldsymbol{\omega}_0, \mathbf{A}_0), \ \ 1 \le n \le N-1;$$

near any given $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{P}_{1,3l}$*:*

$$\begin{aligned}
&\lim_{\mathcal{P}_{1,3}^{-} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_0(\boldsymbol{\omega}, \mathbf{A}) = -\infty, \\
&\lim_{\mathcal{P}_{1,3}^{-} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n-1}(\boldsymbol{\omega}_0, \mathbf{A}_0), \ \ 1 \le n \le N-1, \\
&\lim_{\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3l} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n}(\boldsymbol{\omega}_0, \mathbf{A}_0), \ \ 0 \le n \le N-2, \\
&\lim_{\mathcal{P}_{1,3l}^{+} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_{N-1}(\boldsymbol{\omega}, \mathbf{A}) = +\infty;
\end{aligned}$$

and near any given $(\boldsymbol{\omega}_0, \mathbf{A}_0) \in \mathcal{P}_5$*:*

$$\begin{aligned}
&\lim_{\mathcal{P}_{1,3r}^{+} \cup \mathcal{P}_{1,3r} \cup \mathcal{P}_{1,3}^{-} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_0(\boldsymbol{\omega}, \mathbf{A}) = -\infty, \quad \lim_{\mathcal{P}_{1,3r}^{+} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_1(\boldsymbol{\omega}, \mathbf{A}) = -\infty, \\
&\lim_{\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3l} \cup \mathcal{P}_5 \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_n(\boldsymbol{\omega}_0, \mathbf{A}_0), \ 0 \le n \le N-3, \\
&\lim_{\mathcal{P}_{1,3r} \cup \mathcal{P}_{1,3}^{-} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n-1}(\boldsymbol{\omega}_0, \mathbf{A}_0), \ 1 \le n \le N-2, \\
&\lim_{\mathcal{P}_{1,3r}^{+} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_n(\boldsymbol{\omega}, \mathbf{A}) = \lambda_{n-2}(\boldsymbol{\omega}_0, \mathbf{A}_0), \ 2 \le n \le N-1, \\
&\lim_{\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3l} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_{N-2}(\boldsymbol{\omega}, \mathbf{A}) = +\infty, \quad \lim_{\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3}^{-} \ni (\boldsymbol{\omega},\mathbf{A}) \to (\boldsymbol{\omega}_0,\mathbf{A}_0)} \lambda_{N-1}(\boldsymbol{\omega}, \mathbf{A}) = +\infty.
\end{aligned}$$

And consequently, λ_n *restricted in* $\mathcal{P}_{1,3}^{-} \cup \mathcal{P}_{1,3r}$ *and* $\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3l} \cup \mathcal{P}_5$ *is continuous for each* $0 \le n \le N-3$*, and* λ_{N-2} *restricted in* $\mathcal{P}_{1,3}^{-} \cup \mathcal{P}_{1,3r}$ *and* $\mathcal{P}_{1,3l}^{+} \cup \mathcal{P}_{1,3l}$ *is continuous.*

Theorem 5.4. *Similar results in Theorem* 5.3 *hold for* $\mathcal{O}_{1,3}^{\mathbb{C}}$*,* $\mathcal{P}_{1,3}$*,* $\mathcal{P}_{1,3r}$*,* $\mathcal{P}_{1,3l}$*,* $\mathcal{P}_{1,3}^{-}$*,* $\mathcal{P}_{1,3r}^{+}$*, and* $\mathcal{P}_{1,3l}^{+}$ *replaced by* $\mathcal{O}_{2,3}^{\mathbb{C}}$*,* $\mathcal{P}_{2,3}$*,* $\mathcal{P}_{2,3r}$*,* $\mathcal{P}_{2,3l}$*,* $\mathcal{P}_{2,3}^{-}$*,* $\mathcal{P}_{2,3r}^{+}$*, and* $\mathcal{P}_{2,3l}^{+}$*, separately.*

Combining Theorems 5.1–5.4 yields the continuity and discontinuity sets of the n-th eigenvalue function in $\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}$:

Theorem 5.5. *Each* $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P}_5$ *has exactly* $N-2$ *eigenvalues, and each* $(\boldsymbol{\omega}, \mathbf{A}) \in \mathcal{P} \backslash \mathcal{P}_5$ *has exactly* $N-1$ *eigenvalues, and each* $(\boldsymbol{\omega}, \mathbf{A}) \in \left(\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}\right) \backslash \mathcal{P}$ *has exactly* N *eigenvalues, and* λ_n*,* $0 \le n \le N-1$*, are continuous in* $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}\right) \backslash \mathcal{P}$ *and not continuous at each point of* $\mathcal{P}$*.*

Remark 5.1. If the n-th eigenvalue function λ_n is simple at some $(\boldsymbol{\omega}_0, \mathbf{A}_0)$ in the continuity set $\left(\Omega_N^{\mathbb{R},+} \times \mathcal{B}^{\mathbb{C}}\right) \backslash \mathcal{P}$ for some $0 \le n \le N-1$, the results about differentiability of continuous eigenvalue branches in [22, Theorems 4.2-4.4 and 4.7] can be applied to λ_n in a neighborhood of $(\boldsymbol{\omega}_0, \mathbf{A}_0)$.

References

[1] F. V. Atkinson, Discrete and Continuous Boundary Problems, Academic Press, New York, 1964.

[2] M. Bohner, O. Dosly, The discrete Prüfer transformation, Proc. Amer. Math. Soc. 129 (2001) 2715–2725.

[3] M. Bohner, O. Dosly, W. Kratz, Sturmian and spectral theory for discrete symplectic systems, Trans. Amer. Math. Soc. 361 (2009) 3109–3123.

[4] S. Clark, A spectral analysis for self-adjoint operators generated by a class of second order difference equations, J. Math. Anal. Appl. 197 (1996) 267–285.

[5] S. Clark, F. Gesztesy, W. Renger, Trace formulas and Borg-type theorems for matrix-valued Jacobi and Dirac finite difference operators, J. Differ. Equ. 219 (2005) 144–182.

[6] R. Courant, D. Hilbert, Methods of Mathematical Physics, Interscience Publishers, New York, 1953.

[7] M. Eastham, Q. Kong, H. Wu, A. Zettl, Inequalities among eigenvalues of Sturm-Liouville problems, J. Inequal. Appl. 3 (1999) 25–43.

[8] W. N. Everitt, M. Möller, A. Zettl, Discontinuous dependence of the n-th Sturm-Liouville eigenvalue, Int. Ser. Numer. Math. 123 (1997) 145–150.

[9] A. Jirari, Second-order Sturm-Liouville difference equations and orthogonal polynomials, Mem. Amer. Math. Soc. 113 (1995).

[10] Q. Kong, H. Wu, A. Zettl, Dependence of the n-th Sturm-Liouville eigenvalue on the problem, J. Differ. Equ. 156 (1999) 328–354.

[11] Q. Kong, H. Wu, A. Zettl, Geometric aspects of Sturm-Liouville problems, I. Structures on spaces of boundary conditions, Proc. Roy. Soc. Edinb. Sect. A Math. 130 (2000) 561–589.

[12] H. Lv, Y. Shi, Error estimate of eigenvalues of perturbed second-order discrete Sturm-Liouville problems, Linear Algebra Appl. 430 (2009) 2389–2415.

[13] W. Peng, M. Racovitan, H. Wu, Geometric aspects of Sturm-Liouville problems, V. Natural loops of boundary conditions for monotonicity of eigenvalues and their applications, Pac. J. Appl. Math. 4 (2006) 253–273.

[14] J. Poeschel, E. Trubowitz, Inverse Spectral Theory, Academic Press, New York, 1987.

[15] Y. Shi, Spectral theory for a class of discrete linear Hamiltonian systems, J. Math. Anal. Appl. 289 (2004) 554–570.

[16] Y. Shi, Weyl-Titchmarsh theory for a class of discrete linear Hamiltonian systems, Linear Algebra Appl. 416 (2006) 452–519.

[17] Y. Shi, S. Chen, Spectral theory of second–order vector difference equations, J. Math. Anal. Appl. 239 (1999) 195–212.

[18] Y. Shi, S. Chen, Spectral theory of higher-order discrete vector Sturm-Liouville problems, Linear Algebra Appl. 323 (2001) 7–36.

[19] H. Sun, Y. Shi, Eigenvalues of second-order difference equations with coupled boundary conditions, Linear Algebra Appl. 414 (2006) 361–372.

[20] Y. Wang, Y. Shi, Eigenvalues of second-order difference equations with periodic and

anti-periodic boundary conditions, J. Math. Anal. Appl. 309 (2005) 56–69.

[21] A. Zettl, Sturm-Liouville Theory, Mathematical Surveys Monographs, vol. 121, Amer. Math. Soc., 2005.

[22] H. Zhu, S. Sun, Y. Shi, H. Wu, Dependence of discrete Sturm-Liouville eigenvalues on problems, submitted for publication.